



Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

Marine Ichthyoplankton Data From the Canadian Beaufort Sea Shelf, July and September, 1984

D.B. Chipertzak, G.E. Hopky, M.J. Lawrence
and G. Lacho

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

February 1990

**Canadian Data Report of
Fisheries and Aquatic Sciences
No. 779**



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of Fisheries and Aquatic Sciences

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport statistique canadien des sciences halieutiques et aquatiques

Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Data Report of
Fisheries and Aquatic Sciences 779

February 1990

MARINE ICHTHYOPLANKTON DATA FROM THE CANADIAN BEAUFORT SEA SHELF,
JULY AND SEPTEMBER, 1984

by

D.B. Chipertzak, G.E. Hopky, M.J. Lawrence and G. Lacho

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

This is the 43rd Data Report
from the Central and Arctic Region, Winnipeg

PREFACE

This study was funded by the Northern Oil and Gas Action Program (NOGAP), through the Department of Fisheries and Oceans, Central and Arctic Region. It is one of a series of projects being executed under NOGAP B.2, to provide background data for assessing the implications of hydrocarbon development and production on critical estuarine and marine habitats of the Canadian Arctic Coastal Shelf. This document constitutes NOGAP Report B2.41.

(c) Minister of Supply and Services Canada 1990

Cat. no. FS 97-13/779E

ISSN 0706-6465

Correct citation for this publication is:

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

TABLE OF CONTENTS

	<u>Page</u>
PREFACE	ii
ABSTRACT/RESUME	iv
INTRODUCTION	1
METHODS	1
RESULTS	2
ACKNOWLEDGMENTS	2

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1 Location and detailed map of Beaufort Sea study area	5
2 Stations sampled in July and September, 1984	6

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Summary of plankton stations sampled in the Beaufort Sea, 1984.	7
2 Plankton summary of Wisconsin and neuston tows conducted in the Canadian Beaufort Sea, July and September, 1984	8
3 Stomach fullness codes used in Table 15	9
4 Scientific and common names of fishes captured in the Canadian Beaufort Sea Shelf, 1984	9
5 Number, percent by number, and frequency of occurrence for larval fish captured by Wisconsin tows in the Beaufort Sea Shelf during July and September, 1984.	10
6 Larval fish catch summary for Wisconsin tows on the Beaufort Sea Shelf, 1984	11
7 Larval fish number 100 m ⁻³ for Wisconsin tows on the Beaufort Sea Shelf, 1984	12
8 Number, percent by number, and frequency of occurrence for larval fish captured by neuston tows on the Beaufort Sea Shelf, 1984.	13
9 Larval fish catch summary for neuston tows on the Beaufort Sea Shelf, 1984.	14

<u>Table</u>	<u>Page</u>
10 Larval fish number 100 m ⁻³ for neuston tows on the Beaufort Sea Shelf, 1984	15
11 Standard and total lengths, and wet weight of larval fish captured in the Beaufort Sea, 1984	16
12 Mean length and weight of larval <i>B. saida</i> for July and September, 1984.	30
13 Mean length and weight of larval <i>G. tricuspis</i> for July and September, 1984	30
14 Mean length and weight of larval <i>M. quadricornis</i> for July, 1984.	30
15 Standard length and stomach fullness for larval fish captured in the Beaufort Sea, 1984	31
16 Food item totals for <i>B. saida</i> captured in the Canadian Beaufort Sea Shelf, 1984	34
17 Food item totals for <i>M. quadricornis</i> captured in the Canadian Beaufort Sea Shelf, 1984	40
18 Food item totals for <i>G. tricuspis</i> captured in the Canadian Beaufort Sea Shelf, 1984	41
19 Food item totals for <i>Liparis</i> sp. captured in the Canadian Beaufort Sea Shelf, 1984	42
20 List of comment codes used in Tables 16 to 19	42
21 Frequency of occurrence and percent by number of food items found in <i>B. saida</i> stomachs	43
22 Frequency of occurrence and percent by number of food items found in <i>M. quadricornis</i> stomachs, July 1984	44
23 Frequency of occurrence and percent by number of food items found in <i>G. tricuspis</i> stomachs, July and September 1984	44
24 Frequency of occurrence and percent by number of food items found in <i>Liparis</i> sp. stomachs, July 1984	45
25 Parasites found in larval fish stomachs	45

ABSTRACT

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

Ichthyoplankton were collected from 760 μ m Wisconsin net and 500 μ m neuston net plankton tows in July and September of 1984 along the Canadian Beaufort Sea Shelf. A total of 1408 larval fish representing six families and nine species groups were caught in the 34 plankton tows. Arctic cod (Boreogadus saida) was the most abundant larval fish species caught both in the Wisconsin and neuston tows.

Standard length, total length, and wet-weight measurements were made on 1123 larval fish of which 1097 were Arctic cod. The July mean standard length for Arctic cod was 13.4 mm with a standard deviation of 2.7 mm, and a mean weight of 0.015 g with a standard deviation of 0.010 g. In September, Arctic cod mean standard length was 26.3 mm, standard deviation of 3.8 mm, and the mean weight was 0.130 g with a standard deviation of 0.058 g.

A total of 126 fish were analyzed for stomach contents of which four were found to be empty. Copepods were the most frequently occurring food item in all four species groups analysed with Calanoida being the most frequently occurring and the most abundant food item for both B. saida and the fourhorn sculpin, Myoxcephalus quadricornis.

Nine stomachs, all from B. saida, contained the parasite (Trematoda: Hemiuridae).

Key words: Arctic; coastal waters; ichthyoplankton; larval fishes; Boreogadus saida; Arctic cod; distribution; diet.

RÉSUMÉ

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

En juillet et septembre 1984, on a récolté l'ichthyoplancton à l'aide de filets Wisconsin (760 μm) et de filets à neuston (500 μm) le long de la partie canadienne de la plate-forme de la mer de Beaufort. Trente-quatre traits ont permis de capturer un total de 1408 larves de poisson représentant six familles et neuf groupes d'espèces. Parmi les larves de poisson capturées, la saïda franc (Boreogadus saïda) a été l'espèce la plus abondante et ce, indépendamment du filet utilisé (à neuston ou Wisconsin).

Des mesures de la longueur standard, de la longueur totale et du poids frais ont été effectuées pour 1123 larves de poisson, dont 1097 de saïda franc. En juillet, la longueur standard moyenne des larves de saïda franc était de 13,4 mm, avec un écart-type de 2,7 mm, et le poids moyen était de 0,015 g, avec un écart-type de 0,010 g. en septembre, la longueur standard moyenne des larves de saïda franc était de 26,3 mm, avec un écart-type de 3,8 mm, et le poids moyen était de 0,130 g, avec un écart-type de 0,058 g.

Le contenu stomacal a été analysé chez 126 poissons; quatre de ces poissons avaient l'estomac vide. Les copépodes constituaient l'article alimentaire le plus fréquent dans les quatre groupes d'espèces analysés. Calanoida était l'article alimentaire le plus fréquent et le plus abondant chez B. saïda et chez Myoxcephalus quadricornis (chaboisseau à quatre cornes).

Neuf estomacs, appartenant tous à des poissons de l'espèce B. saïda, étaient parasités (Trematoda: Hemiuridae).

Mots-clés: Arctique; eaux côtières; ichthyoplancton; larves de poisson; Boreogadus saïda; saïda franc; distribution; régime alimentaire.

INTRODUCTION

Ichthyoplankton data presented in this report were collected as part of the Beaufort Shelf Fish Habitat Research Subproject (B.2.1). This Subproject was one component of the Critical Arctic Estuarine and Marine Habitat Project (B.2) of the Northern Oil and Gas Action Program (NOGA-P). The objectives of the subproject were:

1. To conduct research towards identifying, in spatial and temporal terms, areas of significance to marine and estuarine fish species of the Beaufort Sea Shelf (from Herschel Island to Cape Bathurst) (Fig. 1).
2. To characterize these areas of important marine and estuarine habitat in terms of community composition, and in terms of chemical, physical and biological parameters.
3. To describe the feeding habits of selected species of pelagic and demersal fish species of the Beaufort Sea Shelf in relation to habitat and season (open water vs ice covered).

Ichthyoplankton samples were collected for four consecutive summers in the Beaufort Sea beginning in 1984. The data in this report are from 1984. Temperature, salinity and density data for 1984 are reported by Hopky et al. (1986). Stomach data for *Boreogadus saida* (Arctic cod) captured from otter trawl tows done in 1984 are reported by Lacho (1986).

Past studies along the Canadian Beaufort Sea Shelf have had their emphasis on juvenile and adult fish (Galbraith and Hunter 1975), and have been mostly nearshore (Kendel et al. 1975; Percy 1975; Lawrence et al. 1984; Bond and Erickson 1989). The study of larval fish in the Canadian Beaufort Sea has been to date largely ignored, except for Tuktoyaktuk Harbour which has been reported on by Ratynski (1983); yet the larval fish stage is an important time in any fish population because the mortality rate at this stage is so high that even small increases in this rate can severely affect a year class strength (Bagenal and Braum 1978).

Sampling was done on board the chartered M.V. Sequel, a 13 m, wooden hull vessel. Due to the size of the vessel and the nature of its hull, site selection for sampling stations was often at the mercy of ice and weather conditions present at the time of sailing.

METHODS

Coordinates for sampling stations (Table 1) were obtained using a combination of radar (Furuno Model CR240) fixes to shore, bathymetric charts and a Magnavox Model MX4102 satellite navigator with a nominal accuracy of ± 0.1 km.

Plankton samples were collected between July 14 to 25 and September 7 to 9 of 1984 (Table 2). Stations sampled are shown in Fig. 2.

Midwater plankton collections were made using a Wisconsin design plankton net with an opening diameter of 0.5 m and a 763 μ m Nitex mesh net. A General Oceanic flowmeter, model #2030, with a standard rotor was affixed to the centre of the opening. One horizontal tow of 20-40 minutes duration was made at each station. Tow depth was calculated by taking the cosine of the wire angle multiplied by the length of wire spooled out. Ship speed was held at approximately 3 knots for the duration of the tow.

One surface tow using a neuston sampler was made at each station except at stations 84A11 and 84A12 where no surface tows were conducted. The neuston sampler has a square opening of 45 cm x 45 cm with a net constructed of 500 μ m Nitex mesh. A more detailed description of the neuston sampler used is given by Mason and Phillips (1986). A General Oceanic flowmeter, model #2030, with a standard rotor was affixed to the opening of the neuston sampler offset 17 cm vertically from the center of the mouth opening. For volumetric calculations an opening of 45 cm x 35 cm was used, as the top 10 cm of the sampler rode out of the water during the tow periods. The sampler was towed 30 m off the starboard side of the vessel in front of the vessel wake using a polyethylene rope. Ship speed was approximately 4-5 knots for the duration of the tow.

Densities were calculated as the number of fish per 100 m³ of water filtered through either the Wisconsin or neuston nets.

All plankton samples were preserved in a solution of seawater and 5% non-buffered formalin. Larval fish were extracted later in the laboratory, placed in 70% ethanol and identified to the lowest possible taxon.

All larval fish excepting those extracted for stomach analysis or those in too poor a condition for further examination were measured for standard length, total length and wet

weight. Length measurements were made using a Nikon Type 102 dissecting microscope and a small ruler marked in 1 mm graduations. Measurements were made to the nearest ± 0.5 mm. Wet weights were obtained by first blotting the larval fish with a paper towel and then weighing them on a Mettler PC440 digital scale with an accuracy of ± 0.001 g. No adjustments for shrinkage were made in any of the measurements.

Larval fish for stomach analysis were arbitrarily selected to provide a representative cross-section of samples by station and season. These fish were first measured for standard length, rounded to the nearest ± 0.5 mm, using the micrometer eyepiece of a Wild MSA dissecting microscope. The stomach was then removed or in cases where the stomach had not yet fully formed, the entire intestine was taken. A code (Table 3) was assigned to each stomach or intestine to indicate the degree of fullness and the content's state of digestion. Stomach contents were then identified and enumerated. Parasites found in larval fish stomachs were identified and enumerated whenever possible.

RESULTS

A total of 1408 larval fish consisting of nine species groups representing six families (Table 4) were captured from plankton tows in July and September of 1984. Eighteen Wisconsin tows were made (Table 2), capturing 1186 larval fish of which eight species groups in five families were represented. Only 16 neuston tows were made (Table 2), capturing 222 larval fish of which three species groups in three families were represented.

Boreogadus saida was the most abundant species found in Wisconsin tow catches making up 93.0% of the catch in July and 95.8% in September (Table 5). B. saida was also the most frequently occurring species found in Wisconsin tow catches being caught in 66.7% of the catches in July and 100% of the catches in September. All Gadidae captured at station 84B09 (Table 6) were too badly damaged or decomposed to positively identify past family level but are probably B. saida and thus would increase B. saida's percent by number and frequency of occurrence for the month of July. The staghorn sculpin, Gymnocanthus tricuspidis, was the next most abundant species captured in Wisconsin tows making up 2% of the catch in July and 4.2% in September (Table 5). It occurred in 50% of the catches both in July and September. Although six other species were caught intermittently through July, only B. saida and G. tricuspidis were caught in Wisconsin tows during September (Table 6 - "B" stations after 84A12).

Densities (number $\cdot 100 \text{ m}^{-3}$) of fish by species and station for Wisconsin tow catches are given in Table (7).

B. saida was also the most abundant species captured using neuston tows, making up 99.1% of the catch in July and 50% of the catch in September (Table 8). Note that only two fish were caught by the six neuston tows done in September (Table 9 - "B" stations after 84A12). The second fish was an unidentified Cottidae captured at station 84B11. In July, B. saida occurred in 70% of the neuston catches. Only two other species were caught in July by the neuston sampler, one Osmerus mordax and one Myoxcephalus quadricornis both of which were captured at station 84A01 (Table 9).

Densities (number $\cdot 100 \text{ m}^{-3}$) of fish by species and station for all neuston tows are given in Table 10.

Standard length, total length and wet weight measurements were taken on 1123 larval fish (Table 11). Only one specimen each of O. mordax, Anisarchus medius, Triglops pingeli and Aspidophoroides olrikii were measured (Table 11). Virtually all specimens sampled were B. saida, G. tricuspidis or M. quadricornis.

B. saida captured in July had a mean standard length of 13.4 mm with a range of 5.5 to 30.0 mm and a standard deviation of 2.7 while in September the mean standard length was 26.3 mm with a range from 19.5 to 34.0 mm and a standard deviation of 3.8 mm (Table 12). Mean wet weight of B. saida in July was 0.015 g with a standard deviation of 0.010 g while in September the mean wet weight was 0.130 g with a standard deviation of 0.058 g.

G. tricuspidis in July had a mean standard length of 13.6 mm with a range of 11.5 to 15.5 mm and a standard deviation of 1.6 (Table 13). In September G. tricuspidis had a mean standard length of 20.5 mm with a range from 20.0 mm to 21.0 mm and a standard deviation 0.5 mm. The mean wet weight in July was 0.020 g with a standard deviation of 0.009 g and in September the mean wet weight was 0.116 g with a standard deviation of 0.011 g.

Ten M. quadricornis, all in July from station 84A11, were measured (Table 14). The mean standard length was 13.3 mm with a range of 7.0 to 15.5 mm and a standard deviation of 2.3 mm. The mean wet weight was 0.012 g with a range from <0.001 to 0.021 g and a standard deviation of 0.006 g.

A total of 126 fish representing four species groups were examined for their stomach content (Table 15). Of these 99 were B. saida

of which three had empty stomachs. As well, 13 G. tricuspis, six M. quadricornis and four Liparis sp. were examined and all exhibited varying degrees of fullness except one M. quadricornis whose stomach was empty.

A detailed list of food items with totals and comments for the B. saida specimens examined is given in Table 16, for M. quadricornis in Table 17, for G. tricuspis in Table 18 and for Liparis sp. in Table 19. Comment codes are defined in Table 20.

Copepods were the most dominant food group found in the stomachs of all four fish species groups examined (Tables 21 - 24). Calanoids were the most frequently observed food item as well as being the most numerous food item found in both B. saida and M. quadricornis stomachs (Tables 21 and 22).

In B. saida (Table 21), unidentified Calanoida occurred in 83.6% of the stomachs analysed from July and occurred in all stomachs analysed from September. Unidentified Calanoida made up 61.6% by number from July and 63.8% by number from September of food items counted. Pseudocalanus minutus was the next most frequently occurring food item at 26.9% from July as well as the next most numerous food item counted at 31.9% by number, while in September Arcarti clausi occurred slightly more often than P. minutus (59.4% as compared to 56.3%) but P. minutus was the next most abundant species found after Calanoida at 26.2% as compared to A. clausi at 2.2%.

Only copepods were found in the six M. quadricornis stomachs examined (Table 22). Calanoida occurred the most frequently in 50% of the stomachs and was the most abundant food item making up 84.8% by number.

P. minutus was the most frequently occurring food item found in G. tricuspis stomachs, occurring 84.6% of the time, but the diatom Coscinodiscus sp. contributed the greatest percent by number at 49.0% (Table 23). In the four Liparis sp. stomachs the copepod P. minutus occurred the most frequently, being found in 75% of the stomachs analysed and made up the greatest percent by number at 87% (Table 24).

Nine fish, all B. saida, contained the parasite (Trematoda: Hemifuridae) in their stomachs (Table 25).

ACKNOWLEDGMENTS

The authors would like to thank Mr. L. deMarch and Mr. T. Maddeford for field assistance as well as Dr. R. Korczynski who assisted during the September survey. Thanks to Mr. J. Ostrick

and staff at the Inuvik Research Laboratory for logistic support, to Polar Continental Shelf Project for their field operations support and Mr. Ray Ratynski of North South Consultants Ltd. for larval fish identification. Neil Stallard of LGL Ltd. did the stomach content analysis and Margaret Keast provided laboratory assistance; we thank both. The skipper of the M.V. Sequel was L. Pearson. Mr. R. Gervais provided computer assistance and advice. Final manuscript preparation was completed by Barb Hyman and Karla DeCaigny.

REFERENCES

- BAGENAL, T.B., and E. BRAUM. 1978. Eggs and early life history, p. 165-201. In T.B. Bagenal (ed.) Methods for assessment of fish production in fresh waters. Blackwell Scientific Publications, Oxford. 365 p.
- BOND, W.A., and R.N. ERICKSON. 1989. Summer studies of the nearshore fish community at Phillips Bay, Beaufort Sea coast, Yukon. Can. Tech. Rep. Fish. Aquat. Sci. 1676: vi + 102 p.
- GALBRAITH, D.F., and J.G. HUNTER. 1975. Fishes of offshore waters and Tuktoyaktuk vicinity. Beaufort Sea Project, Tech. Rep. 7: 47 p.
- HOPKY, G.E., D.B. CHIPERZAK, and M.J. LAWRENCE. 1986. Seasonal salinity, temperature and density data for the Canadian Beaufort Sea Shelf, 1984-1985. Can. Data Rep. Fish. Aquat. Sci. 593: iv + 249 p.
- KENDEL, R.E., R.A.C. JOHNSTON, U. LOBSIGER, and M.D. KOZAK. 1975. Fishes of the Yukon coast. Beaufort Sea Project, Tech. Rep. 6: 114 p.
- LACHO, G. 1986. Analysis of Arctic cod stomach contents from the Beaufort Shelf, July and August, 1984. Can. Data Rep. Fish. Aquat. Sci. 614: iv + 10 p.
- LAWRENCE, M.J., G. LACHO, and S. DAVIES. 1984. A survey of the coastal fishes of the southeastern Beaufort Sea. Can. Tech. Rep. Fish. Aquat. Sci. 1220: x + 178 p.
- MASON, J.C., and A.C. PHILLIPS. 1986. An improved otter surface sampler. Fish. Bull. 84: 480-484.
- PERCY, R. 1975. Fishes of the outer Mackenzie Delta. Beaufort Sea Project Tech. Rep. 8: 114 p.

RATYNSKI, R.A. 1983. Mid-summer ichthyo-
plankton populations of Tuktoyaktuk
Harbour, N.W.T. Can. Tech. Rep. Fish.
Aquat. Sci. 1218: iv + 21 p.




Fig. 1. Location and detailed map of Beaufort Sea study area.




Fig. 2. Stations sampled in July and September, 1984.

Table 1. Summary of plankton stations sampled in the Beaufort Sea, 1984.

Station	Latitude		Longitude		Date Sampled	Time GMT	Total Depth (m)
	Deg	Min	Deg	Min			
84A01	69	49.5	132	37.0	Jul 14	0100	9
84A02	70	2.0	132	20.0	Jul 15	1625	20
84A03	70	12.4	132	21.6	Jul 15	2045	30
84A04	70	17.3	131	53.5	Jul 15	2300	33
84A05	70	6.4	131	0.5	Jul 16	0600	12
84A06	69	34.1	136	22.5	Jul 23	2000	10
84A07	69	33.2	137	0.1	Jul 23	2335	17
84A08	69	33.6	137	6.4	Jul 23	0220	34
84B09	69	31.7	138	44.6	Jul 24	2000	14
84A10	69	36.5	138	49.0	Jul 24	2330	36
84A11	69	40.0	135	14.0	Jul 25	1200	5
84A12	69	43.5	133	55.0	Jul 25	1650	4
84B01	69	51.8	132	54.0	Sep 7	2300	12
84B02	70	4.0	132	26.0	Sep 7	0335	24
84B03	70	13.0	132	22.0	Sep 8	1900	34
84B04	70	20.0	131	50.0	Sep 8	2325	36
84B05	70	6.0	131	0.0	Sep 9	1640	12
84B11	70	13.0	131	16.0	Sep 9	2100	25

* GMT - Greenwich Mean Time.

Table 2. Plankton summary of Wisconsin and neuston tows conducted in the Canadian Beaufort Sea, July and September, 1984.

Plankton Sample Number	Station	Date	Gear Type	Mesh Size (μ m)	Tow Type	Maximum Tow Depth (m)	Volume Filtered (m^3)
1	84A01	Jul 14	NEUST	500	HOR	0.0	693
2	84A01	Jul 14	WISCN	760	HOR	6.2	531
3	84A02	Jul 15	NEUST	500	HOR	0.0	488
4	84A02	JUL 15	WISCN	760	HOR	13.7	240
5	84A03	Jul 15	WISCN	760	HOR	10.2	427
6	84A03	Jul 15	NEUST	500	HOR	0.0	544
7	84A04	JUL 15	NEUST	500	HOR	0.0	409
8	84A04	Jul 15	WISCN	760	HOR	11.3	370
9	84A05	Jul 16	WISCN	760	HOR	5.1	428
10	84A05	Jul 16	NEUST	500	HOR	0.0	538
11	84A06	Jul 23	NEUST	500	HOR	0.0	515
12	84A06	Jul 23	WISCN	760	HOR	5.1	361
13	84A07	Jul 23	NEUST	500	HOR	0.0	498
14	84A07	Jul 23	WISCN	760	HOR	13.7	448
15	84A08	Jul 23	NEUST	500	HOR	0.0	594
16	84A08	Jul 23	WISCN	760	HOR	11.1	477
17	84B09	Jul 24	NEUST	500	HOR	0.0	321
18	84B09	Jul 24	WISCN	760	HOR	10.2	248
19	84A10	Jul 24	NEUST	500	HOR	0.0	497
20	84A10	Jul 24	WISCN	760	HOR	8.2	383
21	84A11	Jul 25	WISCN	760	HOR	N/A*	N/A
22	84A12	Jul 25	WISCN	760	HOR	4.0	555
29	84B01	Sep 07	NEUST	500	HOR	0.0	532
30	84B01	Sep 07	WISCN	760	HOR	4.7	395
31	84B02	Sep 07	NEUST	500	HOR	0.0	637
32	84B02	Sep 07	WISCN	760	HOR	11.0	483
33	84B03	Sep 08	NEUST	500	HOR	0.0	567
34	84B03	Sep 08	WISCN	760	HOR	11.0	497
35	84B04	Sep 08	NEUST	500	HOR	0.0	496
36	84B04	Sep 08	WISCN	760	HOR	11.7	392
37	84B05	Sep 09	NEUST	500	HOR	0.0	579
38	84B05	Sep 09	WISCN	760	HOR	7.5	442
39	84B11	Sep 09	NEUST	500	HOR	0.0	609
40	84B11	Sep 09	WISCN	760	HOR	12.4	534

* N/A = Not available

Table 3. Stomach fullness codes used in Table 15.

Code	Description
0	Empty
1	1/3 full, digested
2	1/3 full, good condition
3	2/3 full, digested
4	2/3 full, good condition
5	Full, digested
6	Full, good condition

Table 4. Scientific and common names of fishes captured in the Canadian Beaufort Sea Shelf, 1984.

Scientific Name	Common Name
Family Osmeridae <u>Osmerus mordax</u> (Mitchill)	Rainbow smelt
Family Gadidae <u>Boreogadus saida</u> (Lepechin) <u>Eliginus gracilis</u> (Tilesius)	Arctic cod Saffron cod
Family Cottidae <u>Myoxcephalus quadricornis</u> (Linnaeus) <u>Gymnocanthus tricuspis</u> (Reinhardt) <u>Triglops pingeli</u> Reinhardt	Fourhorn sculpin Staghorn sculpin Ribbed sculpin
Family Cyclopteridae <u>Liparis</u> sp.	Snailfish
Family Stichaeidae <u>Anisarchus medius</u> (Reinhardt)	Stout eelblenny
Family Agonidae <u>Aspidophoroides olriki</u> Lutken	Arctic alligatorfish

Table 5. Number, percent by number, and frequency of occurrence for larval fish captured by Wisconsin tows in the Beaufort Sea Shelf during July and September, 1984.

Species	JULY			SEPT		
	Number	%	Freq Occ	Number	%	Freq Occ
Gadidae (unid)	24	2.2	8.3	0	0	0
<u>B. saida</u>	1037	93.0	66.7	68	95.8	100
<u>E. gracilis</u>	1	0.1	8.3	0	0	0
Cottidae (unid)	2	0.2	16.7	0	0	0
<u>G. tricuspis</u>	22	2.0	50.0	3	4.2	50
<u>M. quadricornis</u>	16	1.4	41.7	0	0	0
<u>T. pingell</u>	1	0.1	8.3	0	0	0
<u>A. olriki</u>	1	0.1	8.3	0	0	0
<u>Liparis sp.</u>	8	0.7	33.3	0	0	0
<u>A. medius</u>	1	0.1	8.3	0	0	0
Unid. fish	2	0.2	16.7	0	0	0

Table 6. Larval fish catch summary for Wisconsin tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																							
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	T
<i>O. mordax</i>	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	0
Gadidae (unid)	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	0
<i>B. saia</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	A
<i>E. gracilis</i>	1	2	3	4	5	6	7	8	9	0	1	1	1	1	1	1	1	1	1	1	1	1	1	L
Cottidae (unid)																								
<i>G. tricusps</i>	14	1	3		2			1		1		1		1		1		1		1		1		25
<i>M. quadricornis</i>	2	1						1		2	10													16
<i>T. pingelli</i>																								1
<i>A. olrik</i>																								1
<i>Liparis</i> sp.	3					1	1			3														8
<i>A. medius</i>																								1
Unid. fish	1	1																						2
TOTAL	4	432	17	158	10	316	65	47	24	31	10	1	1	1	27	20	20	1	2	1186				

Table 7. Larval fish number·100 m⁻³ for Wisconsin tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																			
	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	8	8	8	8	8
<i>O. mordax</i>	8	4	4	4	4	4	4	4	4	4	4	4	4	4	4	8	8	8	8	8
Gadidae (unid)	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	8	8	8	8	8
<i>B. saida</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8	8	8	8	8
<i>E. gracilis</i>	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	8	8	8	8	8
Cottidae (unid)																				
<i>G. tricuspid</i>																				
<i>M. quadricornis</i>	0.4	0.4																		
<i>T. pingelli</i>																				
<i>A. olrik</i>																				
<i>Liparis</i> sp.	1.3																			
<i>A. medius</i>	0.4																			
Unid. fish	0.2	0.4	0.7																	
TOTAL	0.8	180.0	4.6	42.6	2.3	87.6	14.5	9.8	9.7	8.1	0.2	0.3	5.6	4.0	5.1	0.2	0.4			

* - Not available

Table 8. Number, percent by number and frequency of occurrence of larval fish captured by neuston tows on the Beaufort Sea Shelf during July and September, 1984.

Species	JULY			SEPT		
	Number	%	Freq Occ	Number	%	Freq Occ
<u>O. mordax</u>	1	0.5	10.0	0	0	0
<u>B. saida</u>	218	99.1	70.0	1	50	16.7
<u>Cottidae (unid.)</u>	0	0.0	0.0	1	50	16.7
<u>M. quadricornis</u>	1	0.5	10.0	0	0	0

Table 9. Larval fish catch summary for neuston tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																							
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
O. mordax	1																							
Gadidae (unid)	1																							
B. saida	5	166	1	2	26	10	8	1																
E. gracilis																								
Cottidae (unid)																								
G. tricusps																								
M. quadricornis	1																							
T. pingelli																								
A. olrik																								
Liparis sp.																								
A. medius																								
Unid. fish																								
TOTAL	2	0	5	166	1	2	26	10	0	8	0	1	0	0	1	0	0	0	1	222				

Table 10. Larval fish number·100 m⁻³ for neuston tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS															
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
O. mordax	0.1															
Gadidae (untd)																
B. saida		0.9	40.6	0.2	0.4	5.2	1.7	1.6		0.2						
E. gracilis																
Cottidae (untd)																
G. tricuspis																
M. quadricornis	0.1															
T. pingelli																
A. olrik																
Liparis sp.																
A. medius																
Unid. fish																
TOTAL	0.2	0.0	0.9	40.6	0.2	0.4	5.2	1.7	0.0	1.6	0.0	0.2	0.0	0.0	0.0	0.2

Table 11. Standard and total lengths, and wet weight of larval fish captured in the Beaufort Sea, 1984.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
1	40	O. mordax	9.0	0.002	4	537	B. salda	14.0	0.017
4	270	B. salda	15.5	0.028	4	538	B. salda	10.0	0.006
4	271	B. salda	13.0	0.015	4	539	B. salda	12.0	0.013
4	273	B. salda	12.0	0.009	4	540	B. salda	12.0	0.013
4	274	B. salda	11.0	0.007	4	541	B. salda	11.5	0.010
4	275	B. salda	15.0	0.023	4	542	B. salda	13.0	0.014
4	277	B. salda	13.0	0.018	4	543	B. salda	13.0	0.012
4	278	B. salda	10.0	0.007	4	544	B. salda	11.5	0.009
4	281	B. salda	13.0	0.015	4	545	B. salda	11.0	0.006
4	282	B. salda	12.5	0.013	4	546	B. salda	11.0	0.007
4	283	B. salda	14.0	0.019	4	547	B. salda	12.0	0.011
4	284	B. salda	13.5	0.021	4	548	B. salda	12.5	0.011
4	285	B. salda	13.0	0.017	4	549	B. salda	11.0	0.011
4	288	B. salda	9.5	0.007	4	550	B. salda	13.0	0.012
4	289	B. salda	12.0	0.006	4	551	B. salda	16.0	0.029
4	291	B. salda	10.0	0.006	4	552	B. salda	13.0	0.014
4	292	B. salda	17.5	0.040	4	553	B. salda	12.0	0.008
4	293	B. salda	13.5	0.019	4	554	B. salda	12.5	0.013
4	294	B. salda	10.0	0.006	4	555	B. salda	11.0	0.007
4	295	B. salda	17.0	0.030	4	556	B. salda	12.0	0.009
4	513	B. salda	16.0	0.025	4	557	B. salda	13.0	0.014
4	514	B. salda	16.0	0.031	4	558	B. salda	14.0	0.018
4	515	B. salda	12.0	0.009	4	559	B. salda	12.0	0.010
4	516	B. salda	14.0	0.024	4	560	B. salda	11.0	0.013
4	517	B. salda	13.5	0.016	4	561	B. salda	12.5	0.015
4	518	B. salda	19.5	0.024	4	562	B. salda	11.0	0.012
4	519	B. salda	5.5	0.007	4	563	B. salda	12.0	0.006
4	520	B. salda	15.5	0.025	4	564	B. salda	15.5	0.028
4	521	B. salda	12.0	0.012	4	565	B. salda	11.5	0.011
4	522	B. salda	11.0	0.010	4	566	B. salda	12.0	0.010
4	523	B. salda	12.0	0.013	4	567	B. salda	12.0	0.011
4	524	B. salda	17.0	0.036	4	568	B. salda	12.0	0.009
4	525	B. salda	14.5	0.019	4	570	B. salda	12.0	0.012
4	526	B. salda	15.0	0.026	4	571	B. salda	10.5	0.008
4	527	B. salda	12.5	0.014	4	572	B. salda	14.5	0.018
4	528	B. salda	14.5	0.022	4	573	B. salda	11.0	0.009
4	529	B. salda	12.0	0.011	4	574	B. salda	12.0	0.009
4	530	B. salda	13.5	0.011	4	575	B. salda	14.0	0.018
4	531	B. salda	12.0	0.010	4	576	B. salda	9.5	0.007
4	532	B. salda	14.0	0.013	4	577	B. salda	11.0	0.005
4	533	B. salda	12.0	0.011	4	578	B. salda	12.0	0.012
4	534	B. salda	14.0	0.020	4	795	B. salda	16.0	0.021
4	535	B. salda	13.0	0.018	4	796	B. salda	16.0	0.022
4	536	B. salda	12.0	0.009	4	797	B. salda	14.5	0.020

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WEI GHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WEI GHT (g)
4	798 B. salda	10.5	11.5	0.009	4	1014 B. salda	14.0	15.0	0.018
4	799 B. salda	14.0	15.0	0.018	4	1015 B. salda	13.0	14.0	0.013
4	800 B. salda	15.0	16.0	0.028	4	1016 B. salda	14.5	15.5	0.022
4	801 B. salda	6.5	7.5	0.009	4	1017 B. salda	14.0	15.0	0.020
4	802 B. salda	10.0	11.0	0.007	4	1018 B. salda	13.5	14.5	0.014
4	803 B. salda	15.5	16.5	0.021	4	1019 B. salda	12.5	13.5	0.014
4	804 B. salda	13.0	14.0	0.014	4	1020 B. salda	12.0	13.0	0.013
4	805 B. salda	12.5	13.5	0.013	4	1021 B. salda	13.5	14.5	0.016
4	806 B. salda	17.0	18.0	0.018	4	1022 B. salda	14.0	15.0	0.019
4	807 B. salda	17.0	18.0	0.012	4	1023 B. salda	12.0	13.0	0.013
4	808 B. salda	14.0	15.0	0.019	4	1024 B. salda	11.0	11.5	0.007
4	809 B. salda	11.5	12.5	0.011	4	1025 B. salda	15.0	16.0	0.022
4	810 B. salda	13.0	14.0	0.016	4	1026 B. salda	13.0	14.0	0.015
4	812 B. salda	13.5	14.5	0.017	4	1027 B. salda	14.5	15.5	0.024
4	813 B. salda	11.0	12.0	0.010	4	1028 B. salda	15.0	16.0	0.022
4	814 B. salda	12.0	13.0	0.010	4	1029 B. salda	11.5	12.5	0.008
4	815 B. salda	14.0	15.0	0.013	4	1030 B. salda	12.0	13.0	0.014
4	816 B. salda	13.5	14.5	0.015	4	1031 B. salda	15.0	16.0	0.020
4	818 B. salda	12.0	13.0	0.013	4	1032 B. salda	14.5	15.5	0.021
4	819 B. salda	14.5	15.5	0.021	4	1033 B. salda	14.0	15.0	0.015
4	820 B. salda	14.5	15.5	0.021	4	1034 B. salda	13.0	14.0	0.017
4	821 B. salda	12.0	13.0	0.014	4	1035 B. salda	15.0	16.0	0.022
4	822 B. salda	10.0	11.0	0.010	4	1036 B. salda	12.5	13.5	0.014
4	823 B. salda	13.0	14.0	0.019	4	1037 B. salda	11.0	12.0	0.010
4	824 B. salda	12.5	13.5	0.012	4	1038 B. salda	10.5	11.0	0.006
4	825 B. salda	13.0	14.0	0.016	4	1039 B. salda	14.0	15.0	0.016
4	826 B. salda	16.0	17.0	0.030	4	1040 B. salda	13.0	14.0	0.013
4	827 B. salda	13.5	14.5	0.018	4	1041 B. salda	12.0	13.0	0.013
4	828 B. salda	11.0	12.0	0.009	4	1042 B. salda	14.0	15.0	0.019
4	829 B. salda	11.0	12.0	0.009	4	1043 B. salda	17.0	18.0	0.011
4	830 B. salda	12.0	13.0	0.012	4	1044 B. salda	13.5	14.5	0.015
4	831 B. salda	14.0	15.0	0.017	4	1045 B. salda	15.0	16.0	0.020
4	832 B. salda	13.0	14.0	0.016	4	1046 B. salda	14.0	15.0	0.018
4	1002 B. salda	14.0	15.0	0.015	4	1047 B. salda	12.5	13.5	0.011
4	1003 B. salda	7.5	8.5	0.011	4	1048 B. salda	12.5	13.5	0.010
4	1004 B. salda	16.0	17.0	0.025	4	1049 B. salda	10.5	11.5	0.006
4	1005 B. salda	15.0	16.0	0.017	4	1050 B. salda	13.0	14.0	0.013
4	1006 B. salda	14.0	15.0	0.016	4	1051 B. salda	12.5	13.5	0.012
4	1007 B. salda	13.5	14.5	0.017	4	1052 B. salda	12.0	13.0	0.008
4	1008 B. salda	14.0	15.0	0.017	4	1053 B. salda	14.5	15.5	0.017
4	1009 B. salda	12.5	13.5	0.014	4	1054 B. salda	16.0	17.0	0.020
4	1010 B. salda	11.5	12.0	0.010	4	1055 B. salda	11.0	12.0	0.008
4	1011 B. salda	13.0	14.0	0.015	4	1056 B. salda	13.5	14.5	0.012
4	1013 B. salda	17.0	18.0	0.028	4	1057 B. salda	15.0	16.0	0.019

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4 1058	B. saida	14.5	15.5	0.017	4 1111	B. saida	15.0	16.0	0.019
4 1059	B. saida	13.0	14.0	0.017	4 1112	B. saida	15.0	16.0	0.021
4 1060	B. saida	12.0	13.0	0.009	4 1113	B. saida	13.5	14.5	0.013
4 1061	B. saida	14.0	15.0	0.020	4 1114	B. saida	11.0	12.0	0.008
4 1062	B. saida	11.0	12.0	0.008	4 1115	B. saida	13.0	14.0	0.014
4 1063	B. saida	13.5	14.5	0.014	4 1116	B. saida	14.5	15.5	0.020
4 1064	B. saida	12.5	13.5	0.013	4 1117	B. saida	13.0	14.0	0.013
4 1065	B. saida	11.0	12.0	0.007	4 1118	B. saida	12.0	13.0	0.013
4 1066	B. saida	14.5	15.5	0.017	4 1119	B. saida	13.5	14.5	0.015
4 1067	B. saida	12.0	13.0	0.013	4 1121	B. saida	12.0	13.0	0.015
4 1068	B. saida	12.5	13.5	0.012	4 1122	B. saida	14.5	15.5	0.014
4 1069	B. saida	12.0	13.0	0.010	4 1123	B. saida	12.0	13.0	0.010
4 1070	B. saida	13.0	14.0	0.013	4 1124	B. saida	13.5	14.5	0.014
4 1071	B. saida	13.0	14.0	0.011	4 1125	B. saida	14.0	15.0	0.020
4 1072	B. saida	16.0	17.0	0.021	4 1126	B. saida	15.0	16.0	0.025
4 1073	B. saida	15.0	16.0	0.028	4 1127	B. saida	15.5	17.0	0.024
4 1074	B. saida	13.0	14.0	0.012	4 1128	B. saida	15.5	16.5	0.019
4 1075	B. saida	12.5	13.5	0.015	4 1129	B. saida	12.0	13.0	0.011
4 1076	B. saida	11.5	12.5	0.012	4 1130	B. saida	13.5	14.5	0.018
4 1077	B. saida	15.0	16.0	0.021	4 1131	B. saida	14.0	15.0	0.020
4 1079	B. saida	12.5	13.5	0.011	4 1132	B. saida	16.0	17.0	0.022
4 1080	B. saida	10.0	11.0	0.009	4 1133	B. saida	11.0	12.0	0.008
4 1081	B. saida	14.0	15.0	0.016	4 1134	B. saida	14.0	15.0	0.014
4 1082	B. saida	15.0	16.0	0.023	4 1135	B. saida	12.5	13.5	0.010
4 1083	B. saida	10.0	11.0	0.008	4 1136	B. saida	13.0	14.0	0.014
4 1084	B. saida	11.0	12.0	0.013	4 1137	B. saida	12.0	13.0	0.013
4 1085	B. saida	13.0	14.0	0.015	4 1138	B. saida	12.0	13.0	0.010
4 1094	B. saida	15.0	16.0	0.023	4 1139	B. saida	12.0	13.0	0.010
4 1095	B. saida	13.0	14.0	0.013	4 1140	B. saida	11.0	12.0	0.007
4 1096	B. saida	14.0	15.0	0.022	4 1141	B. saida	13.5	14.5	0.014
4 1097	B. saida	16.5	17.5	0.026	4 1142	B. saida	11.5	12.5	0.012
4 1098	B. saida	16.5	17.5	0.026	4 1143	B. saida	12.0	13.0	0.013
4 1099	B. saida	13.5	14.5	0.015	4 1144	B. saida	12.5	13.5	0.012
4 1100	B. saida	15.0	16.0	0.023	4 1145	B. saida	14.0	15.0	0.017
4 1101	B. saida	12.0	13.0	0.011	4 1146	B. saida	13.0	14.0	0.013
4 1102	B. saida	15.0	16.0	0.018	4 1147	B. saida	13.5	14.5	0.020
4 1103	B. saida	12.0	13.0	0.014	4 1148	B. saida	13.5	14.5	0.014
4 1104	B. saida	11.5	12.5	0.009	4 1149	B. saida	10.5	11.5	0.010
4 1105	B. saida	15.0	16.0	0.019	4 1150	B. saida	12.0	13.0	0.014
4 1106	B. saida	13.0	14.0	0.016	4 1151	B. saida	15.0	16.0	0.023
4 1107	B. saida	13.0	14.0	0.015	4 1152	B. saida	12.0	13.0	0.009
4 1108	B. saida	14.5	15.5	0.021	4 1153	B. saida	10.0	11.0	0.008
4 1109	B. saida	13.0	14.0	0.019	4 1154	B. saida	12.0	13.0	0.014
4 1110	B. saida	11.5	12.5	0.009	4 1155	B. saida	14.0	15.0	0.017

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WEI T (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WEI T (g)
4	1156 B. salda	12.5	13.5	0.014	4	1202 B. salda	12.0	13.0	0.010
4	1157 B. salda	13.5	14.5	0.012	4	1203 B. salda	14.5	15.5	0.019
4	1158 B. salda	9.0	9.5	0.003	4	1204 B. salda	12.0	13.0	0.009
4	1159 B. salda	15.5	16.5	0.021	4	1205 B. salda	11.0	12.0	0.008
4	1160 B. salda	13.5	14.5	0.018	4	1207 B. salda	13.0	14.0	0.015
4	1161 B. salda	14.0	15.0	0.018	4	1208 B. salda	10.0	10.5	0.005
4	1162 B. salda	12.0	13.0	0.010	4	1210 B. salda	13.0	14.0	0.012
4	1163 B. salda	12.0	13.0	0.010	4	1211 B. salda	10.5	11.5	0.007
4	1164 B. salda	13.0	14.0	0.016	4	1212 B. salda	12.5	13.5	0.011
4	1165 B. salda	13.5	14.5	0.018	4	1213 B. salda	10.5	11.5	0.008
4	1166 B. salda	12.0	13.0	0.014	4	1214 B. salda	12.0	13.0	0.014
4	1167 B. salda	12.5	13.5	0.013	4	1215 B. salda	10.0	11.0	0.006
4	1169 B. salda	9.0	9.5	0.004	4	1216 B. salda	12.0	13.0	0.011
4	1170 B. salda	12.0	13.0	0.012	4	1217 B. salda	11.0	12.0	0.008
4	1171 B. salda	10.0	11.0	0.007	4	1218 B. salda	11.0	12.0	0.007
4	1172 B. salda	16.5	17.5	0.023	4	1219 B. salda	16.0	17.0	0.023
4	1173 B. salda	16.0	17.0	0.028	4	1220 B. salda	13.0	14.0	0.016
4	1174 B. salda	13.5	14.5	0.013	4	1221 B. salda	13.0	14.0	0.014
4	1175 B. salda	10.0	10.5	0.008	4	1222 B. salda	16.0	17.0	0.019
4	1176 B. salda	14.0	15.0	0.016	4	1223 B. salda	13.0	14.0	0.015
4	1177 B. salda	10.0	11.0	0.009	4	1224 B. salda	12.0	13.0	0.011
4	1178 B. salda	18.0	19.0	0.034	4	1225 B. salda	13.0	14.0	0.017
4	1179 B. salda	10.0	11.0	0.008	4	1226 B. salda	13.0	14.0	0.017
4	1180 B. salda	11.0	12.0	0.008	4	1227 B. salda	14.0	15.0	0.021
4	1181 B. salda	12.5	13.5	0.011	4	1228 B. salda	15.0	16.0	0.023
4	1182 B. salda	11.5	12.5	0.009	4	1229 B. salda	15.0	16.0	0.017
4	1183 B. salda	14.0	15.0	0.021	4	1230 B. salda	14.5	15.5	0.020
4	1184 B. salda	10.5	11.5	0.007	4	1231 B. salda	14.0	15.0	0.020
4	1186 B. salda	11.0	12.0	0.006	4	1232 B. salda	12.0	13.0	0.012
4	1187 B. salda	12.0	13.0	0.014	4	1233 B. salda	14.0	15.0	0.020
4	1188 B. salda	13.0	14.0	0.016	4	1234 B. salda	17.5	18.5	0.038
4	1189 B. salda	15.0	16.0	0.023	4	1235 B. salda	13.0	14.0	0.019
4	1190 B. salda	12.5	13.5	0.014	4	1236 B. salda	11.0	12.0	0.010
4	1191 B. salda	11.5	12.5	0.012	4	1237 B. salda	14.5	15.5	0.026
4	1192 B. salda	14.5	15.5	0.024	4	1238 B. salda	12.0	13.0	0.012
4	1193 B. salda	12.5	13.5	0.014	4	1239 B. salda	11.0	11.5	0.008
4	1194 B. salda	13.5	14.5	0.016	4	1240 B. salda	15.0	16.0	0.023
4	1195 B. salda	11.5	12.5	0.007	4	1241 B. salda	16.0	17.0	0.035
4	1196 B. salda	15.5	16.5	0.019	4	1242 B. salda	13.0	14.0	0.018
4	1197 B. salda	12.0	13.0	0.009	4	1243 B. salda	13.5	14.5	0.015
4	1198 B. salda	15.5	16.5	0.022	4	1244 B. salda	12.0	13.0	0.014
4	1199 B. salda	11.5	12.5	0.007	4	1245 B. salda	12.5	13.5	0.013
4	1200 B. salda	13.5	14.5	0.014	4	1246 B. salda	13.0	14.0	0.014
4	1201 B. salda	10.5	11.5	0.008	4	1247 B. salda	12.0	13.0	0.010

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4 1248	B. saida	11.5	12.5	0.011	4	G. tricuspid	15.0	17.5	0.026
4 1249	B. saida	11.5	12.0	0.010	4	G. tricuspid	15.0	17.0	0.033
4 1250	B. saida	14.0	15.0	0.021	4	G. tricuspid	11.5	13.0	0.011
4 1251	B. saida	11.5	12.5	0.013	4	G. tricuspid	13.0	15.0	0.015
4 1252	B. saida	13.5	14.5	0.014	4	G. tricuspid	15.0	17.5	0.030
4 1253	B. saida	10.5	11.0	0.006	4	A. medius	19.5	21.0	0.013
4 1254	B. saida	12.0	13.0	0.014	5	B. saida	12.0	13.0	0.009
4 1255	B. saida	10.0	10.5	0.007	5	B. saida	9.5	10.0	0.002
4 1257	B. saida	9.5	10.0	0.003	5	B. saida	10.0	11.0	0.006
4 1258	B. saida	10.0	10.5	0.005	5	B. saida	12.5	13.5	0.014
4 1284	B. saida	14.0	15.0	0.019	5	B. saida	9.0	10.0	0.003
4 1285	B. saida	12.5	13.5	0.012	5	B. saida	13.0	14.0	0.014
4 1286	B. saida	15.5	17.0	0.031	5	B. saida	10.5	11.0	0.003
4 1287	B. saida	12.0	13.0	0.013	5	B. saida	9.5	10.5	0.003
4 1288	B. saida	13.0	14.0	0.015	5	B. saida	9.5	10.5	0.003
4 1289	B. saida	15.5	16.5	0.021	5	B. saida	10.0	11.0	0.005
4 1290	B. saida	14.5	15.5	0.019	5	B. saida	10.0	11.0	0.005
4 1291	B. saida	17.5	18.5	0.027	5	B. saida	12.0	13.0	0.010
4 1292	B. saida	15.0	16.0	0.023	5	B. saida	8.0	8.5	0.000
4 1294	B. saida	15.5	16.5	0.023	5	B. saida	7.5	8.0	0.000
4 1295	B. saida	18.0	19.0	0.042	5	B. saida	7.5	8.0	0.000
4 1296	B. saida	15.0	16.0	0.023	5	G. tricuspid	11.5	13.0	0.011
4 1297	B. saida	12.0	13.0	0.014	7	B. saida	12.0	13.0	0.009
4 1298	B. saida	16.0	17.0	0.030	7	B. saida	9.5	9.5	0.003
4 1299	B. saida	11.0	12.0	0.011	7	B. saida	11.0	11.0	0.006
4 1300	B. saida	14.0	15.0	0.023	7	B. saida	10.0	10.0	0.005
4 1301	B. saida	17.5	19.0	0.037	7	B. saida	10.0	10.0	0.006
4 1302	B. saida	14.0	15.0	0.020	7	B. saida	9.0	9.0	0.003
4 1303	B. saida	11.0	12.0	0.011	7	B. saida	11.5	12.0	0.007
4 1304	B. saida	15.0	16.0	0.026	7	B. saida	10.5	11.0	0.004
4 1305	B. saida	13.0	14.0	0.021	7	B. saida	10.5	11.0	0.007
4 1306	B. saida	14.0	15.0	0.020	7	B. saida	9.0	9.0	0.002
4 1307	B. saida	15.0	16.0	0.030	7	B. saida	11.0	12.0	0.011
4 1308	B. saida	14.5	15.5	0.025	7	B. saida	11.0	11.0	0.006
4 1309	B. saida	14.0	15.0	0.021	7	B. saida	10.0	10.0	0.005
4 1310	B. saida	15.5	16.5	0.027	7	B. saida	11.0	12.0	0.007
4 1311	B. saida	12.5	13.5	0.011	7	B. saida	11.0	11.0	0.006
4 1312	B. saida	14.5	15.5	0.025	7	B. saida	7.0	7.0	0.002
4 1313	B. saida	14.5	15.5	0.022	7	B. saida	8.5	8.5	0.004
4 1314	B. saida	14.0	15.0	0.022	7	B. saida	9.0	9.5	0.006
4 1315	B. saida	15.0	16.0	0.021	7	B. saida	9.0	9.0	0.005
4 1316	B. saida	14.0	15.0	0.022	7	B. saida	9.0	9.5	0.009
4 1317	B. saida	12.5	13.5	0.016	7	B. saida	14.0	15.0	0.016
4 20	G. tricuspid	15.0	17.5	0.027	7	B. saida	9.5	10.0	0.005

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
7 855	B. salda	9.5	9.5	0.002	7 901	B. salda	8.0	8.0	0.003
7 856	B. salda	10.5	11.0	0.008	7 902	B. salda	9.5	10.0	0.006
7 857	B. salda	10.0	10.0	0.006	7 903	B. salda	10.5	11.0	0.003
7 858	B. salda	10.5	11.0	0.006	7 904	B. salda	9.5	10.0	0.007
7 859	B. salda	10.0	10.5	0.004	7 905	B. salda	8.5	8.5	0.002
7 861	B. salda	11.0	11.5	0.011	7 906	B. salda	9.0	9.0	0.004
7 862	B. salda	10.5	11.0	0.007	7 907	B. salda	9.0	9.0	0.003
7 863	B. salda	10.0	10.5	0.006	7 908	B. salda	12.0	13.0	0.010
7 864	B. salda	10.0	10.5	0.006	7 909	B. salda	10.0	10.0	0.003
7 865	B. salda	9.5	9.5	0.005	7 910	B. salda	12.0	12.5	0.009
7 866	B. salda	10.5	11.0	0.007	7 911	B. salda	9.0	9.0	0.003
7 867	B. salda	10.0	10.0	0.003	7 912	B. salda	9.0	9.5	0.004
7 868	B. salda	9.0	9.0	0.002	7 913	B. salda	9.5	10.0	0.005
7 869	B. salda	9.5	10.0	0.007	7 914	B. salda	9.5	10.0	0.005
7 870	B. salda	8.5	8.5	0.002	7 915	B. salda	9.0	9.0	0.004
7 871	B. salda	9.0	9.0	0.004	7 916	B. salda	9.0	9.0	0.004
7 872	B. salda	11.0	11.0	0.005	7 917	B. salda	9.0	9.5	0.003
7 873	B. salda	9.0	9.0	0.007	7 918	B. salda	10.0	10.5	0.003
7 874	B. salda	9.5	10.0	0.004	7 919	B. salda	9.5	10.0	0.004
7 875	B. salda	9.5	10.0	0.004	7 920	B. salda	9.5	9.5	0.005
7 876	B. salda	10.5	11.0	0.005	7 921	B. salda	9.0	9.5	0.004
7 877	B. salda	9.0	9.0	0.003	7 922	B. salda	8.0	8.0	0.002
7 878	B. salda	9.0	9.0	0.002	7 923	B. salda	9.5	10.0	0.004
7 879	B. salda	10.5	11.0	0.009	7 924	B. salda	8.5	8.5	0.001
7 880	B. salda	9.0	9.0	0.003	7 926	B. salda	11.0	11.5	0.008
7 881	B. salda	10.0	10.0	0.004	7 927	B. salda	10.0	10.5	0.007
7 882	B. salda	11.0	11.5	0.008	7 928	B. salda	9.0	9.5	0.005
7 883	B. salda	10.0	10.5	0.007	7 930	B. salda	11.5	12.5	0.011
7 884	B. salda	10.0	10.5	0.004	7 931	B. salda	9.0	9.5	0.004
7 885	B. salda	9.0	9.0	0.003	7 933	B. salda	11.0	11.5	0.005
7 886	B. salda	10.0	10.5	0.005	7 934	B. salda	7.5	7.5	0.000
7 887	B. salda	8.5	9.0	0.004	7 935	B. salda	11.0	12.0	0.010
7 888	B. salda	9.5	10.0	0.006	7 936	B. salda	9.5	10.0	0.005
7 889	B. salda	10.5	11.0	0.008	7 937	B. salda	11.0	12.0	0.009
7 890	B. salda	9.0	9.0	0.004	7 938	B. salda	10.0	10.5	0.006
7 891	B. salda	9.5	10.0	0.004	7 939	B. salda	8.0	8.0	0.002
7 893	B. salda	9.5	10.0	0.003	7 940	B. salda	10.5	11.0	0.009
7 894	B. salda	13.0	14.0	0.013	7 941	B. salda	10.5	11.0	0.008
7 895	B. salda	12.0	13.0	0.010	7 942	B. salda	10.0	10.0	0.002
7 896	B. salda	10.0	10.5	0.006	7 943	B. salda	9.5	10.0	0.005
7 897	B. salda	8.0	8.0	0.002	7 944	B. salda	10.0	10.5	0.006
7 898	B. salda	8.0	8.0	0.002	7 945	B. salda	11.0	12.0	0.010
7 899	B. salda	9.5	10.0	0.008	7 946	B. salda	9.5	10.0	0.004
7 900	B. salda	8.0	8.0	0.002	7 947	B. salda	12.0	13.0	0.011

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)
7 948	B. salda	9.0	9.5	0.003	7 994	B. salda	9.5	10.0	0.003
7 949	B. salda	10.0	10.5	0.004	7 995	B. salda	9.5	10.0	0.004
7 950	B. salda	9.5	10.0	0.005	7 996	B. salda	8.0	8.0	0.000
7 951	B. salda	9.0	9.0	0.004	7 997	B. salda	9.0	9.0	0.002
7 952	B. salda	10.5	11.0	0.007	8 47	B. salda	13.5	14.0	0.011
7 953	B. salda	9.0	9.5	0.005	8 48	B. salda	14.0	15.0	0.011
7 954	B. salda	8.0	8.0	0.001	8 49	B. salda	12.5	13.5	0.008
7 955	B. salda	9.0	9.5	0.004	8 50	B. salda	15.0	16.0	0.016
7 956	B. salda	9.0	9.0	0.005	8 52	B. salda	14.0	15.0	0.017
7 957	B. salda	8.0	8.5	0.002	8 53	B. salda	11.0	12.0	0.011
7 958	B. salda	12.0	13.0	0.011	8 54	B. salda	15.0	16.0	0.016
7 959	B. salda	9.0	9.0	0.003	8 55	B. salda	14.0	15.0	0.016
7 960	B. salda	9.0	9.5	0.003	8 56	B. salda	11.5	12.5	0.010
7 961	B. salda	10.0	10.0	0.002	8 57	B. salda	10.5	10.5	0.006
7 962	B. salda	10.5	11.0	0.005	8 58	B. salda	14.0	15.0	0.013
7 963	B. salda	9.5	10.0	0.007	8 59	B. salda	12.5	13.5	0.009
7 964	B. salda	9.5	10.0	0.006	8 60	B. salda	12.0	13.0	0.009
7 965	B. salda	10.0	10.5	0.004	8 61	B. salda	14.0	15.0	0.012
7 966	B. salda	10.5	11.0	0.005	8 62	B. salda	14.0	15.0	0.015
7 967	B. salda	9.0	9.0	0.004	8 63	B. salda	12.0	12.5	0.007
7 968	B. salda	9.5	10.0	0.005	8 64	B. salda	12.5	13.5	0.010
7 969	B. salda	10.0	10.0	0.003	8 65	B. salda	14.0	15.0	0.012
7 970	B. salda	10.0	10.0	0.004	8 66	B. salda	12.0	13.0	0.007
7 971	B. salda	9.0	9.0	0.002	8 67	B. salda	11.5	12.0	0.008
7 972	B. salda	10.0	10.5	0.005	8 68	B. salda	12.0	13.0	0.010
7 973	B. salda	10.0	10.5	0.005	8 69	B. salda	11.5	12.5	0.008
7 974	B. salda	9.5	10.0	0.004	8 70	B. salda	13.0	14.0	0.009
7 975	B. salda	9.0	9.0	0.002	8 71	B. salda	12.0	13.0	0.008
7 976	B. salda	11.5	12.5	0.010	8 72	B. salda	11.5	12.5	0.009
7 977	B. salda	8.0	8.0	0.002	8 73	B. salda	14.0	15.0	0.016
7 978	B. salda	8.5	8.5	0.001	8 74	B. salda	13.5	14.0	0.013
7 979	B. salda	12.0	13.0	0.011	8 75	B. salda	13.0	14.0	0.013
7 980	B. salda	8.0	8.0	0.002	8 76	B. salda	13.0	14.0	0.012
7 981	B. salda	9.0	9.0	0.003	8 77	B. salda	15.0	16.0	0.021
7 982	B. salda	10.0	11.0	0.006	8 78	B. salda	12.0	13.0	0.009
7 983	B. salda	12.0	13.0	0.009	8 79	B. salda	14.0	15.0	0.017
7 984	B. salda	11.0	12.0	0.008	8 80	B. salda	14.0	15.0	0.016
7 985	B. salda	10.5	11.0	0.007	8 81	B. salda	12.0	13.0	0.012
7 986	B. salda	11.5	12.0	0.008	8 82	B. salda	12.0	13.0	0.007
7 987	B. salda	10.0	10.5	0.006	8 83	B. salda	13.5	14.0	0.008
7 988	B. salda	10.0	10.5	0.005	8 84	B. salda	12.0	13.0	0.009
7 989	B. salda	9.5	9.5	0.006	8 85	B. salda	13.0	14.0	0.010
7 990	B. salda	9.5	9.5	0.003	8 86	B. salda	13.0	14.0	0.011
7 991	B. salda	9.5	10.0	0.004	8 87	B. salda	13.0	14.0	0.013
7 992	B. salda	9.5	10.0	0.004					
7 993	B. salda	9.5	10.0	0.004					

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
88	B. saida	13.5	14.5	0.010	140	B. saida	12.0	13.0	0.008
89	B. saida	12.0	13.0	0.010	141	B. saida	11.5	12.5	0.008
90	B. saida	14.0	15.0	0.015	143	B. saida	10.0	10.5	0.008
91	B. saida	13.0	14.0	0.012	144	B. saida	10.5	11.0	0.007
92	B. saida	12.5	13.5	0.009	145	B. saida	9.5	10.0	0.004
93	B. saida	10.0	10.5	0.006	146	B. saida	11.5	12.5	0.008
94	B. saida	15.5	16.5	0.018	147	B. saida	11.0	12.0	0.007
95	B. saida	13.5	14.5	0.011	148	B. saida	13.5	14.5	0.014
97	B. saida	12.5	13.5	0.010	149	B. saida	12.5	14.0	0.013
99	B. saida	11.5	12.5	0.009	150	B. saida	11.0	12.0	0.006
100	B. saida	12.0	13.0	0.010	151	B. saida	10.5	11.5	0.009
101	B. saida	11.0	12.0	0.008	152	B. saida	12.0	13.0	0.008
102	B. saida	13.0	14.0	0.011	153	B. saida	12.5	14.0	0.013
103	B. saida	12.5	13.5	0.010	154	B. saida	13.0	14.0	0.015
104	B. saida	13.0	14.0	0.014	155	B. saida	10.5	11.5	0.009
105	B. saida	13.0	14.0	0.013	156	B. saida	14.0	15.0	0.017
106	B. saida	11.5	12.5	0.011	157	B. saida	11.0	12.0	0.007
107	B. saida	12.5	13.5	0.008	158	B. saida	14.0	15.0	0.013
109	B. saida	9.0	10.0	0.004	159	B. saida	13.5	14.5	0.013
111	B. saida	12.0	13.0	0.011	160	B. saida	10.0	11.0	0.005
112	B. saida	13.0	14.0	0.012	161	B. saida	12.5	13.5	0.011
113	B. saida	12.5	13.5	0.010	162	B. saida	14.0	15.0	0.017
114	B. saida	16.0	17.5	0.024	163	B. saida	9.0	10.0	0.004
116	B. saida	12.0	13.0	0.010	164	B. saida	12.0	13.0	0.011
117	B. saida	14.0	15.0	0.013	165	B. saida	12.0	13.0	0.008
120	B. saida	12.0	13.0	0.008	166	B. saida	13.0	14.0	0.012
121	B. saida	13.0	14.0	0.015	167	B. saida	10.5	11.5	0.007
122	B. saida	13.5	14.5	0.013	169	B. saida	11.0	12.0	0.005
123	B. saida	12.0	13.0	0.009	170	B. saida	10.0	10.5	0.004
124	B. saida	15.5	17.0	0.020	171	B. saida	12.0	13.0	0.007
126	B. saida	12.0	13.0	0.009	172	B. saida	11.0	12.0	0.008
127	B. saida	12.0	13.0	0.011	173	B. saida	11.0	12.0	0.007
128	B. saida	14.0	15.0	0.013	174	B. saida	13.0	14.0	0.010
129	B. saida	13.5	15.0	0.015	175	B. saida	11.5	12.5	0.010
130	B. saida	11.0	12.0	0.006	176	B. saida	11.5	12.5	0.007
131	B. saida	11.0	12.0	0.007	178	B. saida	10.5	11.5	0.007
132	B. saida	13.0	14.0	0.013	179	B. saida	11.0	12.0	0.006
133	B. saida	13.0	14.0	0.012	182	B. saida	11.0	12.0	0.008
134	B. saida	12.0	13.0	0.007	183	B. saida	15.0	16.0	0.017
135	B. saida	13.0	14.0	0.011	184	B. saida	12.0	13.0	0.009
136	B. saida	14.0	15.0	0.013	186	B. saida	11.0	12.0	0.008
137	B. saida	14.0	15.0	0.012	188	B. saida	13.5	14.5	0.013
138	B. saida	12.0	13.0	0.009	189	B. saida	8.0	9.0	0.003
139	B. saida	10.5	11.0	0.004	190	B. saida	12.0	13.0	0.009

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
6	191	B. salda	12.0	0.008	12	335	B. salda	17.0	0.022
8	192	B. salda	10.5	0.005	12	337	B. salda	14.0	0.010
8	193	B. salda	12.0	0.008	12	338	B. salda	17.0	0.024
8	194	B. salda	10.5	0.005	12	339	B. salda	17.5	0.024
8	197	B. salda	6.0	0.002	12	341	B. salda	15.0	0.014
8	198	B. salda	8.5	0.003	12	342	B. salda	17.0	0.020
8	1329	G. trieuapia	11.5	0.006	12	343	B. salda	13.0	0.013
8	1330	G. trieuapia	15.5	0.024	12	344	B. salda	19.0	0.036
8	1331	G. trieuapia	13.0	0.015	12	345	B. salda	17.5	0.024
8	1337	A. olrikii	8.5	0.001	12	346	B. salda	17.0	0.020
10	605	B. salda	30.0	0.188	12	347	B. salda	16.5	0.018
11	1000	B. salda	17.0	0.021	12	348	B. salda	15.0	0.015
11	1001	B. salda	15.5	0.017	12	349	B. salda	14.5	0.014
12	296	B. salda	19.0	0.035	12	350	B. salda	15.0	0.014
12	297	B. salda	16.5	0.023	12	351	B. salda	15.0	0.014
12	298	B. salda	19.0	0.030	12	352	B. salda	12.0	0.009
12	299	B. salda	17.0	0.023	12	354	B. salda	17.5	0.024
12	301	B. salda	16.5	0.017	12	355	B. salda	15.5	0.017
12	302	B. salda	17.0	0.025	12	356	B. salda	14.0	0.014
12	303	B. salda	15.0	0.019	12	357	B. salda	18.0	0.027
12	304	B. salda	16.0	0.017	12	358	B. salda	14.0	0.011
12	305	B. salda	16.0	0.017	12	359	B. salda	16.5	0.026
12	306	B. salda	16.0	0.019	12	360	B. salda	15.0	0.017
12	307	B. salda	16.0	0.019	12	361	B. salda	18.0	0.026
12	308	B. salda	17.0	0.020	12	362	B. salda	15.0	0.017
12	310	B. salda	13.0	0.015	12	363	B. salda	16.5	0.020
12	311	B. salda	16.5	0.023	12	365	B. salda	18.0	0.028
12	312	B. salda	13.0	0.010	12	366	B. salda	16.0	0.021
12	313	B. salda	18.0	0.032	12	368	B. salda	13.0	0.008
12	314	B. salda	19.0	0.039	12	369	B. salda	14.0	0.014
12	316	B. salda	15.0	0.015	12	370	B. salda	18.0	0.026
12	318	B. salda	17.0	0.024	12	371	B. salda	15.0	0.019
12	319	B. salda	17.0	0.026	12	373	B. salda	16.0	0.019
12	321	B. salda	12.5	0.008	12	374	B. salda	16.5	0.018
12	322	B. salda	17.5	0.026	12	375	B. salda	14.0	0.012
12	323	B. salda	11.5	0.020	12	376	B. salda	12.0	0.007
12	324	B. salda	14.0	0.014	12	377	B. salda	15.0	0.013
12	325	B. salda	17.5	0.030	12	378	B. salda	16.0	0.024
12	326	B. salda	17.0	0.028	12	379	B. salda	16.0	0.020
12	328	B. salda	13.5	0.010	12	380	B. salda	14.0	0.010
12	329	B. salda	16.0	0.019	12	381	B. salda	17.0	0.026
12	330	B. salda	16.0	0.018	12	382	B. salda	17.0	0.021
12	331	B. salda	16.0	0.023	12	383	B. salda	15.0	0.015
12	332	B. salda	18.0	0.034	12	384	B. salda	16.0	0.022

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12 385	B. salda	13.0	14.0	0.011	12 616	B. salda	10.0	11.0	0.011
12 387	B. salda	17.0	18.0	0.021	12 617	B. salda	15.0	16.0	0.017
12 388	B. salda	18.0	19.5	0.035	12 618	B. salda	14.0	15.0	0.013
12 389	B. salda	17.0	18.5	0.023	12 620	B. salda	18.5	20.0	0.035
12 390	B. salda	15.0	16.0	0.014	12 621	B. salda	17.5	18.5	0.026
12 393	B. salda	15.5	16.5	0.016	12 622	B. salda	15.0	16.0	0.016
12 394	B. salda	16.0	17.0	0.016	12 623	B. salda	16.5	18.0	0.031
12 396	B. salda	18.0	19.0	0.026	12 624	B. salda	15.0	16.0	0.015
12 397	B. salda	19.0	20.5	0.027	12 625	B. salda	14.0	15.0	0.009
12 398	B. salda	15.0	16.0	0.016	12 628	B. salda	16.0	17.0	0.016
12 399	B. salda	19.0	20.5	0.033	12 629	B. salda	17.5	19.0	0.029
12 401	B. salda	16.5	17.5	0.020	12 630	B. salda	14.5	15.5	0.016
12 402	B. salda	15.0	16.0	0.010	12 632	B. salda	13.5	14.5	0.009
12 403	B. salda	16.5	17.5	0.019	12 633	B. salda	14.0	15.0	0.012
12 404	B. salda	17.0	18.0	0.018	12 634	B. salda	17.0	18.0	0.022
12 405	B. salda	14.5	15.5	0.017	12 635	B. salda	13.0	14.0	0.010
12 406	B. salda	14.0	15.0	0.009	12 636	B. salda	16.5	17.5	0.020
12 407	B. salda	16.5	17.5	0.025	12 637	B. salda	18.0	19.0	0.026
12 408	B. salda	17.5	19.0	0.029	12 638	B. salda	15.5	16.5	0.017
12 409	B. salda	17.0	18.5	0.022	12 639	B. salda	15.0	16.0	0.016
12 410	B. salda	18.0	19.5	0.026	12 640	B. salda	13.5	14.5	0.015
12 411	B. salda	14.0	15.0	0.012	12 641	B. salda	14.5	15.5	0.017
12 412	B. salda	13.0	14.0	0.010	12 642	B. salda	19.0	20.5	0.031
12 413	B. salda	15.0	16.0	0.015	12 643	B. salda	16.5	17.5	0.021
12 414	B. salda	13.0	14.0	0.009	12 644	B. salda	18.0	19.0	0.029
12 415	B. salda	17.0	18.0	0.022	12 645	B. salda	17.0	18.5	0.026
12 416	B. salda	14.5	15.5	0.015	12 646	B. salda	16.0	17.0	0.022
12 417	B. salda	17.0	18.5	0.021	12 647	B. salda	17.0	18.0	0.024
12 418	B. salda	18.5	20.0	0.031	12 648	B. salda	16.0	17.0	0.017
12 419	B. salda	12.0	13.0	0.008	12 649	B. salda	15.0	16.0	0.020
12 422	B. salda	15.0	16.0	0.016	12 651	B. salda	16.0	17.0	0.020
12 423	B. salda	16.0	17.0	0.025	12 652	B. salda	18.5	20.0	0.033
12 426	B. salda	15.0	16.0	0.015	12 653	B. salda	18.0	19.0	0.032
12 427	B. salda	15.0	16.0	0.014	12 654	B. salda	13.5	14.5	0.010
12 606	B. salda	14.0	15.0	0.016	12 655	B. salda	14.0	15.0	0.012
12 607	B. salda	15.0	16.0	0.015	12 656	B. salda	16.5	17.5	0.022
12 608	B. salda	15.0	16.0	0.015	12 657	B. salda	16.0	17.0	0.025
12 609	B. salda	18.0	20.0	0.038	12 658	B. salda	13.0	14.0	0.012
12 610	B. salda	13.0	14.0	0.015	12 659	B. salda	19.0	20.0	0.038
12 611	B. salda	17.0	18.0	0.017	12 660	B. salda	16.0	17.0	0.024
12 612	B. salda	18.0	19.0	0.026	12 661	B. salda	15.0	16.0	0.011
12 613	B. salda	18.0	20.0	0.037	12 662	B. salda	19.0	20.5	0.040
12 614	B. salda	16.5	17.5	0.026	12 663	B. salda	16.0	17.0	0.021
12 615	B. salda	9.5	10.5	0.017	12 664	B. salda	16.5	17.5	0.020

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12 665	B. salda	16.0	17.0	0.021	12 713	B. salda	14.0	15.0	0.013
12 667	B. salda	14.5	15.5	0.015	12 714	B. salda	15.0	16.0	0.016
12 668	B. salda	16.0	17.0	0.014	12 715	B. salda	13.0	14.0	0.013
12 669	B. salda	15.0	16.0	0.012	12 716	B. salda	18.5	20.0	0.034
12 670	B. salda	14.0	15.0	0.016	12 717	B. salda	15.0	16.0	0.019
12 671	B. salda	18.0	19.5	0.031	12 718	B. salda	16.0	17.5	0.023
12 672	B. salda	13.5	14.5	0.013	12 719	B. salda	15.0	16.0	0.013
12 673	B. salda	15.0	16.0	0.015	12 720	B. salda	15.5	16.5	0.016
12 674	B. salda	13.0	14.0	0.014	12 721	B. salda	18.0	19.0	0.032
12 675	B. salda	16.5	18.0	0.024	12 722	B. salda	18.0	19.0	0.027
12 676	B. salda	14.0	15.0	0.009	12 723	B. salda	20.5	22.0	0.045
12 677	B. salda	16.0	17.0	0.021	12 724	B. salda	18.0	19.5	0.032
12 678	B. salda	14.0	15.0	0.018	12 725	B. salda	18.5	19.5	0.030
12 680	B. salda	16.0	17.0	0.021	12 726	B. salda	18.5	20.0	0.034
12 681	B. salda	15.5	16.5	0.018	12 727	B. salda	15.0	16.0	0.016
12 682	B. salda	15.5	16.5	0.016	12 729	B. salda	17.0	18.0	0.021
12 683	B. salda	18.5	20.0	0.039	12 730	B. salda	15.0	16.0	0.015
12 684	B. salda	18.5	20.0	0.035	12 731	B. salda	14.0	15.0	0.018
12 685	B. salda	20.0	22.0	0.041	12 732	B. salda	14.0	15.0	0.016
12 686	B. salda	14.0	15.0	0.015	12 733	B. salda	16.0	17.0	0.024
12 687	B. salda	17.0	18.0	0.032	12 734	B. salda	16.0	17.0	0.021
12 688	B. salda	16.0	17.0	0.020	12 736	B. salda	16.5	17.5	0.019
12 689	B. salda	15.0	16.0	0.018	12 737	B. salda	15.5	17.0	0.019
12 690	B. salda	15.5	16.5	0.016	12 738	B. salda	14.5	15.5	0.015
12 691	B. salda	12.5	13.5	0.008	12 739	B. salda	14.5	15.5	0.020
12 692	B. salda	17.0	18.0	0.027	12 740	B. salda	17.5	19.0	0.027
12 693	B. salda	16.5	18.0	0.020	12 741	B. salda	14.0	15.0	0.014
12 694	B. salda	14.5	15.5	0.014	12 743	B. salda	16.0	17.0	0.018
12 695	B. salda	14.5	15.5	0.015	12 744	B. salda	16.5	17.5	0.020
12 696	B. salda	14.0	15.0	0.014	12 745	B. salda	17.0	18.0	0.021
12 697	B. salda	18.5	19.5	0.031	12 746	B. salda	16.0	17.0	0.023
12 698	B. salda	13.5	14.5	0.012	12 747	B. salda	15.5	16.5	0.016
12 699	B. salda	12.0	13.0	0.007	12 748	B. salda	18.0	19.5	0.031
12 700	B. salda	15.0	16.0	0.014	12 749	B. salda	15.5	16.5	0.017
12 701	B. salda	13.5	14.5	0.013	12 750	B. salda	20.0	20.0	0.032
12 702	B. salda	17.0	18.5	0.028	12 752	B. salda	14.0	15.0	0.016
12 703	B. salda	15.5	16.5	0.016	12 753	B. salda	19.5	21.0	0.042
12 704	B. salda	16.0	17.0	0.018	12 754	B. salda	17.0	18.0	0.022
12 705	B. salda	18.0	20.0	0.030	12 755	B. salda	17.0	18.0	0.032
12 707	B. salda	17.5	18.5	0.020	12 756	B. salda	13.5	14.5	0.007
12 708	B. salda	17.0	18.0	0.023	12 757	B. salda	17.0	18.0	0.022
12 709	B. salda	14.0	15.0	0.009	12 758	B. salda	14.0	15.0	0.015
12 710	B. salda	18.0	19.5	0.031	12 759	B. salda	15.5	16.5	0.018
12 711	B. salda	19.0	20.5	0.030	12 761	B. salda	18.5	20.0	0.029

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)
12	762 B. salda	15.5	16.5	0.019	14	462 B. salda	15.0	16.0	0.017
12	763 B. salda	15.5	16.5	0.016	14	463 B. salda	16.0	17.0	0.025
12	764 B. salda	18.5	20.0	0.030	14	464 B. salda	14.0	15.0	0.017
12	765 B. salda	16.0	17.0	0.016	14	465 B. salda	17.0	18.0	0.029
12	766 B. salda	16.0	17.0	0.018	14	466 B. salda	14.5	15.5	0.013
12	767 B. salda	15.0	16.0	0.017	14	467 B. salda	17.5	19.0	0.030
12	768 B. salda	17.5	19.0	0.028	14	468 B. salda	15.5	16.5	0.024
12	769 B. salda	19.0	20.5	0.036	14	469 B. salda	17.0	18.0	0.034
12	770 B. salda	18.0	19.5	0.036	14	470 B. salda	16.5	17.5	0.021
12	772 B. salda	16.0	17.0	0.018	14	471 B. salda	12.0	13.0	0.010
12	773 B. salda	14.0	15.0	0.017	14	472 B. salda	16.0	17.0	0.017
12	774 B. salda	17.0	18.0	0.024	14	473 B. salda	18.0	19.5	0.031
12	775 B. salda	16.0	17.0	0.020	14	474 B. salda	14.5	15.5	0.020
12	776 B. salda	13.5	14.5	0.011	14	475 B. salda	15.5	16.5	0.023
12	778 B. salda	16.0	17.0	0.018	14	477 B. salda	16.0	17.0	0.025
12	779 B. salda	17.0	18.0	0.020	14	478 B. salda	16.0	17.0	0.024
12	780 B. salda	14.5	15.5	0.013	14	479 B. salda	15.0	16.0	0.019
12	781 B. salda	16.0	17.0	0.019	14	480 B. salda	19.0	21.0	0.037
12	783 B. salda	16.0	17.0	0.025	14	481 B. salda	18.5	20.0	0.040
12	784 B. salda	16.0	17.0	0.021	14	483 B. salda	16.0	17.0	0.021
12	785 B. salda	14.0	15.0	0.015	14	484 B. salda	12.5	13.5	0.012
12	786 B. salda	17.0	18.0	0.026	14	485 B. salda	17.0	18.0	0.029
13	1259 B. salda	14.5	15.5	0.013	14	486 B. salda	15.0	16.0	0.024
13	1260 B. salda	14.5	15.5	0.018	14	487 B. salda	17.5	19.0	0.031
13	1261 B. salda	15.0	16.0	0.014	14	488 B. salda	14.0	15.0	0.015
13	1262 B. salda	17.5	18.5	0.022	14	489 B. salda	15.5	16.5	0.016
13	1263 B. salda	15.0	16.0	0.014	14	490 B. salda	16.5	17.5	0.024
13	1266 B. salda	16.0	17.0	0.020	14	491 B. salda	15.0	16.0	0.022
13	1268 B. salda	16.0	17.0	0.016	14	492 B. salda	14.0	15.0	0.015
13	1270 B. salda	16.0	17.0	0.016	14	493 B. salda	14.5	15.5	0.020
13	1281 B. salda	14.5	15.5	0.014	14	494 B. salda	17.0	18.5	0.028
14	449 B. salda	16.0	17.0	0.013	14	495 B. salda	12.0	13.0	0.008
14	450 B. salda	16.0	17.0	0.024	14	496 B. salda	17.0	19.5	0.031
14	451 B. salda	14.0	15.0	0.018	14	497 B. salda	13.5	14.5	0.016
14	452 B. salda	15.0	16.0	0.020	14	498 B. salda	16.0	17.0	0.022
14	453 B. salda	15.5	16.5	0.022	14	499 B. salda	13.0	14.0	0.012
14	454 B. salda	16.5	17.5	0.021	14	500 B. salda	18.0	19.0	0.038
14	455 B. salda	17.0	18.0	0.026	14	501 B. salda	14.0	15.0	0.014
14	456 B. salda	14.0	15.0	0.012	14	502 B. salda	16.0	17.0	0.028
14	457 B. salda	15.0	16.0	0.014	14	503 B. salda	12.0	13.0	0.008
14	458 B. salda	16.5	17.5	0.033	14	504 B. salda	19.0	20.5	0.039
14	459 B. salda	15.0	16.0	0.021	14	505 B. salda	11.0	12.0	0.007
14	460 B. salda	16.0	17.0	0.022	14	506 B. salda	16.0	17.0	0.022
14	461 B. salda	15.0	16.0	0.016	14	507 B. salda	15.0	16.0	0.021

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	NET WEIGHT (g)
12	762 B. salda	15.5	16.5	0.019	14	462 B. salda	15.0	16.0	0.017
12	763 B. salda	15.5	16.5	0.016	14	463 B. salda	16.0	17.0	0.025
12	764 B. salda	18.5	20.0	0.030	14	464 B. salda	14.0	15.0	0.017
12	765 B. salda	16.0	17.0	0.016	14	465 B. salda	17.0	18.0	0.029
12	766 B. salda	16.0	17.0	0.018	14	466 B. salda	14.5	15.5	0.013
12	767 B. salda	15.0	16.0	0.017	14	467 B. salda	17.5	19.0	0.030
12	768 B. salda	17.5	19.0	0.028	14	468 B. salda	15.5	16.5	0.024
12	769 B. salda	19.0	20.5	0.036	14	469 B. salda	17.0	18.0	0.034
12	770 B. salda	18.0	19.5	0.036	14	470 B. salda	16.5	17.5	0.021
12	772 B. salda	16.0	17.0	0.018	14	471 B. salda	12.0	13.0	0.010
12	773 B. salda	14.0	15.0	0.017	14	472 B. salda	16.0	17.0	0.017
12	774 B. salda	17.0	18.0	0.024	14	473 B. salda	18.0	19.5	0.031
12	775 B. salda	16.0	17.0	0.020	14	474 B. salda	14.5	15.5	0.020
12	776 B. salda	13.5	14.5	0.011	14	475 B. salda	15.5	16.5	0.023
12	778 B. salda	16.0	17.0	0.018	14	477 B. salda	16.0	17.0	0.025
12	779 B. salda	17.0	18.0	0.020	14	478 B. salda	16.0	17.0	0.024
12	780 B. salda	14.5	15.5	0.019	14	479 B. salda	15.0	16.0	0.019
12	781 B. salda	16.0	17.0	0.013	14	480 B. salda	19.0	21.0	0.037
12	783 B. salda	16.0	17.0	0.025	14	481 B. salda	18.5	20.0	0.040
12	784 B. salda	16.0	17.0	0.021	14	483 B. salda	16.0	17.0	0.021
12	785 B. salda	14.0	15.0	0.015	14	484 B. salda	12.5	13.5	0.012
12	786 B. salda	17.0	18.0	0.026	14	485 B. salda	17.0	18.0	0.029
13	1259 B. salda	14.5	15.5	0.013	14	486 B. salda	15.0	16.0	0.024
13	1260 B. salda	14.5	15.5	0.018	14	487 B. salda	17.5	19.0	0.031
13	1261 B. salda	15.0	16.0	0.014	14	488 B. salda	14.0	15.0	0.015
13	1262 B. salda	17.5	18.5	0.022	14	489 B. salda	15.5	16.5	0.016
13	1263 B. salda	15.0	16.0	0.014	14	490 B. salda	16.5	17.5	0.024
13	1266 B. salda	16.0	17.0	0.020	14	491 B. salda	15.0	16.0	0.022
13	1268 B. salda	16.0	17.0	0.016	14	492 B. salda	14.0	15.0	0.015
13	1270 B. salda	16.0	17.0	0.016	14	493 B. salda	14.5	15.5	0.020
13	1281 B. salda	14.5	15.5	0.014	14	494 B. salda	17.0	18.0	0.028
14	449 B. salda	16.0	17.0	0.013	14	495 B. salda	12.0	13.0	0.008
14	450 B. salda	16.0	17.0	0.024	14	496 B. salda	17.0	19.5	0.031
14	451 B. salda	14.0	15.0	0.018	14	497 B. salda	13.5	14.5	0.016
14	452 B. salda	15.0	16.0	0.020	14	498 B. salda	16.0	17.0	0.022
14	453 B. salda	15.5	16.5	0.022	14	499 B. salda	13.0	14.0	0.012
14	454 B. salda	16.5	17.5	0.021	14	500 B. salda	18.0	19.0	0.038
14	455 B. salda	17.0	18.0	0.026	14	501 B. salda	14.0	15.0	0.014
14	456 B. salda	14.0	15.0	0.012	14	502 B. salda	16.0	17.0	0.028
14	457 B. salda	15.0	16.0	0.014	14	503 B. salda	12.0	13.0	0.008
14	458 B. salda	16.5	17.5	0.033	14	504 B. salda	19.0	20.5	0.039
14	459 B. salda	15.0	16.0	0.021	14	505 B. salda	11.0	12.0	0.007
14	460 B. salda	16.0	17.0	0.022	14	506 B. salda	16.0	17.0	0.022
14	461 B. salda	15.0	16.0	0.016	14	507 B. salda	15.0	16.0	0.021

Table 11. Continued.

PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
16	45 B. saia	9.5	9.5	0.000	32	601 B. saia	30.0	32.5	0.170
16	46 B. saia	12.5	13.5	0.007	32	602 B. saia	34.0	37.0	0.277
16	1338 T. pingelli	20.5	23.0	0.062	32	603 B. saia	22.5	24.5	0.071
20	220 B. saia	13.0	13.0	0.006	32	604 B. saia	25.0	27.0	0.094
20	221 B. saia	17.0	18.5	0.031	32	1318 G. tricuspid	20.0	25.0	0.105
20	222 B. saia	17.0	18.5	0.026	34	430 B. saia	29.0	32.0	0.129
20	223 B. saia	18.0	19.0	0.021	34	431 B. saia	31.0	34.0	0.221
20	225 B. saia	15.0	16.0	0.012	34	433 B. saia	25.0	27.0	0.086
20	228 B. saia	17.0	18.0	0.021	34	438 B. saia	22.0	24.0	0.066
20	229 B. saia	16.5	17.5	0.017	34	439 B. saia	26.5	29.5	0.112
20	230 B. saia	13.0	13.0	0.008	34	440 B. saia	27.0	29.5	0.123
20	233 B. saia	13.0	13.0	0.004	34	441 B. saia	31.0	34.0	0.194
20	1086 B. saia	18.0	19.0	0.023	34	442 B. saia	22.5	25.0	0.080
20	1087 B. saia	13.5	14.5	0.012	34	447 B. saia	21.5	23.5	0.059
20	1088 B. saia	14.0	15.0	0.015	36	199 B. saia	28.0	31.0	0.155
20	1090 B. saia	13.5	14.5	0.012	36	205 B. saia	22.0	24.0	0.093
20	1091 B. saia	16.0	17.0	0.021	36	206 B. saia	29.0	32.0	0.172
20	1092 B. saia	15.5	16.5	0.021	36	208 B. saia	26.0	28.0	0.120
20	1093 B. saia	15.0	16.0	0.015	36	210 B. saia	20.0	22.0	0.062
21	2 M. quadricornis	15.0	17.5	0.019	36	211 B. saia	26.0	28.0	0.128
21	3 M. quadricornis	14.5	16.0	0.016	36	212 B. saia	26.0	28.0	0.121
21	4 M. quadricornis	15.5	17.5	0.021	36	214 B. saia	19.5	21.5	0.050
21	5 M. quadricornis	14.0	16.0	0.013	36	217 B. saia	20.0	22.0	0.061
21	6 M. quadricornis	14.0	16.0	0.011	40	1336 G. tricuspid	21.0	24.5	0.127
21	7 M. quadricornis	12.5	14.0	0.009					
21	8 M. quadricornis	14.0	16.0	0.014					
21	9 M. quadricornis	13.5	15.0	0.011					
21	10 M. quadricornis	13.0	14.0	0.005					
21	11 M. quadricornis	7.0	7.0	0.000					
29	219 B. saia	31.0	34.0	0.187					
31	235 B. saia	23.0	25.0	0.071					
32	580 B. saia	27.0	29.0	0.122					
32	582 B. saia	27.0	29.0	0.154					
32	583 B. saia	28.0	30.0	0.147					
32	584 B. saia	22.0	24.0	0.068					
32	585 B. saia	33.0	37.0	0.270					
32	586 B. saia	24.0	26.0	0.089					
32	590 B. saia	27.0	30.0	0.137					
32	593 B. saia	30.5	33.5	0.218					
32	594 B. saia	24.0	26.0	0.104					
32	595 B. saia	25.0	27.0	0.121					
32	598 B. saia	30.0	33.0	0.188					
32	600 B. saia	30.5	33.0	0.168					

Table 12. Mean length and weight of larval B. salda for July and September, 1984.

	JULY				SEPT			
	Mean	Range	Std Dev	N	Mean	Range	Std Dev	N
Std length	13.4	5.5-30.0	2.7	1061	26.3	19.5-34.0	3.8	36
Total length	14.3	6.5-32.5	3.0	1061	28.7	21.5-37.0	4.2	36
Wet weight	0.015	0.007-0.188	0.010	1061	0.130	0.050-0.277	0.058	36

Table 13. Mean length and weight of larval G. tricuspis for July and September, 1984.

	JULY				SEPT			
	Mean	Range	Std Dev	N	Mean	Range	Std Dev	N
Std length	13.6	11.5-15.5	1.6	10	20.5	20.0-21.0	0.5	2
Total length	15.5	12.5-17.5	2.0	10	24.8	24.5-25.0	0.25	2
Wet weight	0.020	0.006-0.033	0.009	10	0.116	0.105-0.127	0.011	2

Table 14. Mean length and weight of larval M. quadricornis for July, 1984.

	JULY			
	Mean	Range	Std Dev	N
Std length	13.3	7.0-15.5	2.26	10
Total length	14.9	7.0-17.5	2.87	10
Wet weight	0.119	<0.001-0.021	0.006	10

Table 15. Standard length and stomach fullness for larval fish captured in the Beaufort Sea, 1984.

Plankton Sample Number	Fish Sam No	Species	Std Length (mm)	Stomach Taken	Stomach Fullness
1	1681	<u>M. quadricornis</u>	11.0	W	1
2	1682	<u>M. quadricornis</u>	10.0	W	0
2	1683	<u>M. quadricornis</u>	11.0	W	3
4	1501	<u>B. saida</u>	10.5	W	3
4	1502	<u>B. saida</u>	11.5	W	5
4	1503	<u>B. saida</u>	14.5	W	5
4	1504	<u>B. saida</u>	14.0	W	5
4	1505	<u>B. saida</u>	11.5	W	3
4	1506	<u>B. saida</u>	11.0	W	3
4	1507	<u>B. saida</u>	14.0	W	5
4	1508	<u>B. saida</u>	11.5	W	1
4	1509	<u>B. saida</u>	15.5	S	6
4	1669	<u>G. tricuspis</u>	11.5	W	5
4	1670	<u>G. tricuspis</u>	11.5	W	3
4	1671	<u>G. tricuspis</u>	11.0	W	3
4	1672	<u>G. tricuspis</u>	13.5	W	3
4	1673	<u>G. tricuspis</u>	14.0	W	3
4	1674	<u>G. tricuspis</u>	14.5	W	5
4	1684	<u>G. tricuspis</u>	13.0	W	3
4	1689	<u>Liparis sp.</u>	20.5	S	6
4	1690	<u>Liparis sp.</u>	13.0	W	5
4	1691	<u>Liparis sp.</u>	9.0	W	3
5	1675	<u>G. tricuspis</u>	11.0	W	3
6	1528	<u>B. saida</u>	8.5	W	1
6	1529	<u>B. saida</u>	7.5	W	3
6	1530	<u>B. saida</u>	9.0	W	3
6	1531	<u>B. saida</u>	8.5	W	3
6	1532	<u>B. saida</u>	8.5	W	3
8	1533	<u>B. saida</u>	11.5	W	5
8	1534	<u>B. saida</u>	12.5	W	5
8	1535	<u>B. saida</u>	13.0	W	5
8	1536	<u>B. saida</u>	14.0	W	5
8	1537	<u>B. saida</u>	14.5	S	6
8	1538	<u>B. saida</u>	14.5	S	4
8	1539	<u>B. saida</u>	15.0	S	6
8	1540	<u>B. saida</u>	14.5	S	4
8	1541	<u>B. saida</u>	15.0	S	4
8	1542	<u>B. saida</u>	16.0	S	4
9	1543	<u>B. saida</u>	*	W	5
9	1544	<u>B. saida</u>	10.5	W	3
9	1545	<u>B. saida</u>	9.0	W	0
9	1546	<u>B. saida</u>	10.0	W	1
9	1547	<u>B. saida</u>	9.5	W	5

Table 15. Continued.

Plankton Sample Number	Fish Sam No	Species	Std Length (mm)	Stomach Taken	Stomach Fullness
9	1548	<u>B. saida</u>	9.5	W	3
9	1549	<u>B. saida</u>	10.5	W	3
9	1550	<u>B. saida</u>	11.5	W	3
9	1551	<u>B. saida</u>	10.5	W	5
9	1552	<u>B. saida</u>	12.5	W	3
12	1553	<u>B. saida</u>	15.0	W	3
12	1554	<u>B. saida</u>	17.0	W	0
12	1555	<u>B. saida</u>	16.0	W	3
12	1556	<u>B. saida</u>	16.0	W	3
12	1557	<u>B. saida</u>	16.5	W	1
12	1558	<u>B. saida</u>	18.5	S	4
12	1559	<u>B. saida</u>	17.0	S	4
12	1560	<u>B. saida</u>	17.5	S	0
12	1561	<u>B. saida</u>	17.0	S	2
12	1562	<u>B. saida</u>	18.5	S	4
12	1676	<u>G. tricuspis</u>	12.0	W	1
12	1677	<u>G. tricuspis</u>	11.0	W	1
13	1563	<u>B. saida</u>	15.5	W	4
13	1565	<u>B. saida</u>	15.5	W	1
13	1567	<u>B. saida</u>	9.0	W	1
13	1569	<u>B. saida</u>	12.5	W	1
13	1570	<u>B. saida</u>	12.5	W	3
13	1572	<u>B. saida</u>	14.0	W	1
13	1574	<u>B. saida</u>	15.5	W	3
13	1577	<u>B. saida</u>	14.5	W	3
13	1581	<u>B. saida</u>	17.0	W	5
13	1583	<u>B. saida</u>	15.5	W	1
14	1692	<u>Liparis sp.</u>	10.5	W	3
15	1589	<u>B. saida</u>	10.0	W	5
15	1590	<u>B. saida</u>	11.0	W	5
15	1591	<u>B. saida</u>	11.0	W	5
15	1592	<u>B. saida</u>	11.5	W	5
15	1593	<u>B. saida</u>	11.5	W	5
15	1594	<u>B. saida</u>	13.0	W	5
15	1595	<u>B. saida</u>	13.5	W	3
15	1596	<u>B. saida</u>	15.0	W	5
16	1678	<u>G. tricuspis</u>	14.0	W	1
16	1688	<u>M. quadricornis</u>	11.5	W	1
19	1597	<u>B. saida</u>	12.5	W	3
19	1598	<u>B. saida</u>	13.0	W	3
19	1599	<u>B. saida</u>	14.0	W	3
19	1600	<u>B. saida</u>	14.5	W	5
19	1601	<u>B. saida</u>	14.5	W	3

Table 15. Continued.

Plankton Sample Number	Fish Sam No	Species	Std Length (mm)	Stomach Taken	Stomach Fullness
19	1602	<u>B. saida</u>	16.0	W	6
19	1603	<u>B. saida</u>	16.5	S	2
19	1604	<u>B. saida</u>	17.0	S	3
20	1679	<u>G. tricusps</u>	14.0	W	3
20	1686	<u>M. quadricornis</u>	11.5	W	1
20	1687	<u>M. quadricornis</u>	12.0	W	1
21	1685	<u>M. quadricornis</u>	15.5	W	1
32	1605	<u>B. saida</u>	11.0	W	3
32	1608	<u>B. saida</u>	24.5	S	6
32	1610	<u>B. saida</u>	25.5	S	4
32	1613	<u>B. saida</u>	25.5	S	6
32	1615	<u>B. saida</u>	23.5	S	6
32	1617	<u>B. saida</u>	26.0	S	5
32	1619	<u>B. saida</u>	28.5	S	3
32	1621	<u>B. saida</u>	*	S	4
32	1623	<u>B. saida</u>	30.0	S	6
32	1627	<u>B. saida</u>	34.0	S	2
34	1628	<u>B. saida</u>	18.5	S	1
34	1630	<u>B. saida</u>	21.0	S	4
34	1632	<u>B. saida</u>	25.0	S	5
34	1634	<u>B. saida</u>	20.5	S	4
34	1636	<u>B. saida</u>	24.5	S	4
34	1638	<u>B. saida</u>	25.5	S	6
34	1639	<u>B. saida</u>	26.0	S	2
34	1641	<u>B. saida</u>	38.5	S	6
34	1645	<u>B. saida</u>	28.5	S	2
34	1647	<u>B. saida</u>	32.0	S	4
36	1648	<u>B. saida</u>	21.0	S	5
36	1650	<u>B. saida</u>	21.0	S	4
36	1652	<u>B. saida</u>	22.0	S	6
36	1654	<u>B. saida</u>	21.0	S	6
36	1656	<u>B. saida</u>	26.5	S	4
36	1658	<u>B. saida</u>	25.0	S	4
36	1660	<u>B. saida</u>	26.0	S	4
36	1662	<u>B. saida</u>	27.5	S	4
36	1664	<u>B. saida</u>	28.5	S	4
36	1666	<u>B. saida</u>	30.0	S	4
36	1680	<u>G. tricusps</u>	*	S	3
38	1667	<u>B. saida</u>	20.0	S	1
40	1668	<u>B. saida</u>	24.0	S	2

* - No measurement taken

W - Intestine (stomach not yet developed)

S - Stomach

Table 16. *Continued.*[illegible]

Table 16. Continui.

[illegible]

Table 16. Continued.

QC = comment code
YL = total

Table 17. Food item totals for M. quadricornis captured in the Canadian Beaufort Sea Shelf, 1984.

Planton Sam. No.	Fish Sam. No.	<u>Limnocalanus macrurus</u>		<u>Calanoida</u>		<u>Cyclopoida</u>	
		CC	TL	CC	TL	CC	TL
1	1681					N2	3
2	1683			N1 6			
				N3 19			
16	1688	Ps	1				
20	1686			Ps 2			
20	1687			Ps 1			
21	1685	Ps	1				

CC = comment code

TL = total

Table 18. Food item totals for *G. tricuspid* captured in the Canadian Beaufort Sea Shelf, 1984.

Plank- ton Fish SAM No.		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL		CC		TL	
-------------------------------------	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--

Table 19. Food item totals for Liparis sp. captured in the Canadian Beaufort Sea Shelf, 1984.

Plankton Sam. No.	Fish Sam. No.	Polychaete (Unid.)		Oncaea borealis		Pseudocalanus minutus		Apherusa glacialis		Gammarus sp.	
		CC	TL	CC	TL	CC	TL	CC	TL	CC	TL
4	1689					C2	1	8		Ju	1
4	1690					Cf	14				
						Ce	125				
4	1691	Ju	1	Cm	1	C5	1				
14	1692	Ju	10								

CC = comment code

TL = total

Table 20. List of comment codes used in Tables 16 to 19.

Code	Comment
Ps	poor condition
P1	Polychaete larvae
La	larva - no specific taxon
Na	nauplii - no specific taxon
Ju	juvenile - no specific taxon
C1	copepodite I
C2	copepodite II
C3	copepodite III
C4	copepodite IV
C5	copepodite V
Cf	adult female (copepodite VI)
Cm	adult male (copepodite VI)
N1	nauplius < 200um
N2	nauplius 200 - 400um
N3	nauplius > 400um
C7	copepodite < 400um (Cyclopoids and Harpacticoids only)
C8	copepodite 400 - 800um (Cyclopoids and Harpacticoids only)
Ce	copepod eggs

Table 21. Frequency of occurrence and percent by number of food items found in B. saida stomachs.

Food Item		July		Sept	
		Freq Occur	% Num	Freq Occur	% Num
Diatoms:	<u>Coscinodiscus sp.</u>	1.5	0.1	12.5	0.2
	<u>Peridinium sp.</u>			9.4	0.1
	<u>Ceratium sp.</u>	1.5	0.1		
Rotifers:	Unid.	3.0	0.1	34.4	0.6
Polychaetes:	Unid.			9.4	0.2
Copepoda:	<u>Acartia clausi</u>			59.4	2.2
	<u>Calanus glacialis</u>	3.0	0.1	46.9	1.0
	<u>Calanus hyperboreus</u>			3.1	<0.1
	<u>Calanus sp.</u>			3.1	<0.1
	<u>Eurytemora sp.</u>			3.1	<0.1
	<u>Derjuginia tolli</u>	1.5	0.2		
	<u>Limnocalanus macrurus</u>	17.9	1.1	3.1	<0.1
	<u>Microcalanus pygmaeus</u>			3.1	<0.1
	<u>Microsetella rosea</u>			9.4	0.1
	<u>Oithona similis</u>	17.9	2.8	34.4	0.7
	<u>Oncaea borealis</u>	4.5	0.2	43.8	0.5
	<u>Pseudocalanus minutus</u>	26.9	31.9	56.3	26.2
	Calanoida (Unid.)	83.6	61.6	100.0	63.8
	Cyclopoida (Unid.)	6.0	0.3	21.9	0.3
Amphipoda:	<u>Apherusa glacialis</u>			3.1	0.1
Gastropoda:	<u>Clione limacina</u>			3.1	<0.1
	<u>Limacina helicina</u>			3.1	<0.1
Bivalvia:	Unid.	9.0	0.5	15.6	0.2
Larvacea:	<u>Oikopleura vanhoeffeni</u>	6.0	1.0	40.6	1.6
	<u>Fritillaria borealis</u>			18.8	0.9
Eggs:	Unid.	3.0	0.2	50.0	1.3

Table 22. Frequency of occurrence and percent by number of food items found in M. quadricornis stomachs, July, 1984.

Food Item		Freq Occur	% Num
Copepoda:	<u>Limnocalanus macrurus</u>	33.3	6.1
	Calanoida (Unid.)	50.0	84.8
	Cyclopoida (Unid.)	16.7	9.1

Table 23. Frequency of occurrence and percent by number of food items found in G. tricuspis stomachs, July and September, 1984.

Food Item		Freq Occur	% Num
Diatoms:	<u>Coscinodiscus</u> sp.	61.5	49.0
	<u>Peridinium</u> sp.	7.7	0.3
	Unid.	7.7	1.7
Polychaetes:	Unid.	15.4	0.6
Copepoda:	<u>Cyclopina</u> sp.	15.4	10.1
	<u>Limnocalanus macrurus</u>	15.4	0.8
	<u>Microsetella rosea</u>	7.7	0.3
	<u>Oithona similis</u>	7.7	0.3
	<u>Oncaea borealis</u>	7.7	0.3
	<u>Pseudocalanus minutus</u>	84.6	20.8
	Calanoida	15.4	0.8
	Harpacticoida	7.7	0.3
Cirripedia:	Unid.	38.5	1.4
Amphipoda:	<u>Apherusa glacialis</u>	7.7	0.6
Bivalvia:	Unid.	23.1	3.4
Stelleroide:	Unid.	7.7	1.1
Larvacea:	<u>Oikopleura vanhoeffeni</u>	7.7	2.8
Eggs:	Unid.	30.8	5.6

Table 24. Frequency of occurrence and percent by number of food items found in Liparis sp. stomachs, July, 1984.

Food Item		Freq Occur	% Num
Polychaetes:	Unid.	50.0	6.8
Copepoda:	<u>Oncaea borealis</u>	25.0	0.6
	<u>Pseudocalanus minutus</u>	75.0	87.0
Amphipoda:	<u>Gammarus</u> sp.	25.0	0.6
	<u>Apherusa glacialis</u>	25.0	4.9

Table 25. Parasites found in larval fish stomachs.

Fish Sam No	Species	Station	Parasite	Count
1507	<u>B. saida</u>	84A02	Trematoda: Hemiuridae	1
1542	<u>B. saida</u>	84A04	Trematoda: Hemiuridae	2
1608	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	2
1619	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1621	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	2
1623	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1627	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1647	<u>B. saida</u>	84B03	Trematoda: Hemiuridae	1
1660	<u>B. saida</u>	84B04	Trematoda: Hemiuridae	1